

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.


**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 5/065, 5/087, 5/107, A61K 38/05, 38/06, 38/07, C07C 257/18, C07D 209/20, 317/60, 333/24, 207/14, 211/26, 207/16, 233/54, 211/30, 211/22, 211/34, 217/26, 295/18, 295/22, 277/28, 249/08	A1	(11) International Publication Number: WO 99/11658 (43) International Publication Date: 11 March 1999 (11.03.99)
(21) International Application Number: PCT/GB98/02605 (22) International Filing Date: 28 August 1998 (28.08.98) (30) Priority Data: 9718392.5 29 August 1997 (29.08.97) GB 9803173.5 13 February 1998 (13.02.98) GB (71) Applicant (for all designated States except US): PROTEUS MOLECULAR DESIGN LTD. [GB/GB]; Beechfield House, Lyme Green Business Park, Macclesfield, Cheshire SK11 0JL (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): LIEBESCHUETZ, John, Walter [GB/GB]; Beechfield House, Lyme Green Business Park, Macclesfield, Cheshire SK11 0JL (GB). WYLIE, William, Alexander [GB/GB]; Beechfield House, Lyme Green Business Park, Macclesfield, Cheshire SK11 0JL (GB). WASZKOWYCZ, Bohdan [GB/GB]; Beechfield House, Lyme Green Business Park, Macclesfield, Cheshire SK11 0JL (GB). MURRAY, Christopher, William [GB/GB]; Beechfield House, Lyme Green Business		Park, Macclesfield, Cheshire SK11 0JL (GB). RIMMER, Andrew, David [GB/GB]; Beechfield House, Lyme Green Business Park, Macclesfield, Cheshire SK11 0JL (GB). WELSH, Pauline, Mary [GB/GB]; Beechfield House, Lyme Green Business Park, Macclesfield, Cheshire SK11 0JL (GB). JONES, Stuart, Donald [GB/GB]; Beechfield House, Lyme Green Business Park, Macclesfield, Cheshire SK11 0JL (GB). ROSCOE, Jonathan, Michael, Ernest [GB/GB]; Beechfield House, Lyme Green Business Park, Macclesfield, Cheshire SK11 0JL (GB). YOUNG, Stephen, Clinton [GB/GB]; Beechfield House, Lyme Green Business Park, Macclesfield, Cheshire SK11 0JL (GB). MORGAN, Phillip, John [GB/GB]; Beechfield House, Lyme Green Business Park, Macclesfield, Cheshire SK11 0JL (GB). (74) Agents: COCKBAIN, Julian et al.; Frank B. Dehn & Co., 179 Queen Victoria Street, London EC4V 4EL (GB). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report.
(54) Title: META-BENZAMIDINE DERIVATIVES AS SERIN PROTEASE INHIBITORS (57) Abstract The invention relates to serine protease inhibitor compound of formula (I), where R ₁ and R ₂ each independently is hydrogen or hydroxyl, alkoxyl, alkyl, aminoalkyl, hydroxyalkyl, alkoxyalkyl, alkoxyacarbonyl, acyloxymethoxycarbonyl or alkylamino optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl; each R ₃ independently is R ₁ , R ₂ , amino, halo, cyano, nitro, thiol, alkylthio, alkylsulfonyl, alkylsulphenyl, alkylsulphonamido, alkylaminosulphonyl, haloalkoxy and haloalkyl; each X independently is C, N, O or S atom or a CO, CR ₁ , C(R ₁) ₂ or NR ₁ group, at least one X being C, CO, CR ₁ or C(R ₁) ₂ , with the proviso that if the benzamidine group is unsubstituted and the X-X group is -CH ₂ C(R ₁) ₂ -, then R ₁ is hydrogen or is attached to the alkylene carbon atom by a heteroatom; L is an organic linker group containing 1 to 5 backbone atoms selected from C, N, O and S, or a branched alkyl or cyclic group; Y is a nitrogen atom or a CR ₁ group or Y and L taken together form a cyclic group; Cy is a saturated or unsaturated, mono or polycyclic, homo or heterocyclic group optionally substituted by groups R ₃ or phenyl optionally substituted by R ₃ ; Lp is a lipophilic organic group, selected from an alkyl, heterocyclic, alkenyl, alkaryl, cycloalkyl, polycycloalkyl, cycloalkenyl, aryl, aralkyl or haloalkyl group or a combination of two or more such groups optionally substituted by one or more of oxa, oxo, aza, thio, halo, amino, hydroxy or R ₃ groups; D is a hydrogen bond donor group; and n is 0, 1 or 2 and salts thereof.		



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

META-BENZAMIDINE DERIVATIVES AS SERIN PROTEASE INHIBITORS

5 This invention relates to compounds which are inhibitors of serine proteases and to pharmaceutical compositions thereof and their use in the treatment of the human or animal body.

The serine proteases are a group of proteolytic enzymes which have a common catalytic mechanism
10 characterized by a particularly reactive Ser residue. Examples of serine proteases include trypsin, tryptase, chymotrypsin, elastase, thrombin, plasmin, kallikrein, Complement C1, acrosomal protease, lysosomal protease, cocoonase, α -lytic protease, protease A, protease B,
15 serine carboxypeptidase II, subtilisin, urokinase, Factor VIIa, Factor IXa, and Factor Xa. The serine proteases have been investigated extensively over a period of several decades and the therapeutic value of inhibitors of serine proteases is well understood.

20 Serine protease inhibitors play a central role in the regulation of a wide variety of physiological process including coagulation, fibrinolysis, fertilization, development, malignancy, neuromuscular patterning and inflammation. It is well known that
25 these compounds inhibit a variety of circulating proteases as well as proteases that are activated or released in tissue. It is also becoming clear that serine protease inhibitors inhibit critical cellular processes, such as adhesion, migration, free radical
30 production and apoptosis. In addition, animal experiments indicate that intravenously administered serine protease inhibitors, variants or cells expressing serine protease inhibitors, provide a protective effect against tissue damage.

35 Serine protease inhibitors have also been predicted to have potential beneficial uses in the treatment of disease in a wide variety of clinical areas such as

- 2 -

oncology, neurology, haematology, pulmonary medicine, immunology, inflammation and infectious disease.

In particular serine protease inhibitors may be beneficial in the treatment of thrombotic diseases, asthma, emphysema, cirrhosis, arthritis, carcinoma, melanoma, restenosis, atheroma, trauma, shock and reperfusion injury.

Thus for example an inhibitor of Factor Xa has value as a therapeutic agent as an anticoagulant, e.g. in the treatment and prevention of thrombotic disorders. The use of a Factor Xa inhibitor as an anticoagulant is desirable in view of the selectivity of its effect. Many clinically approved anticoagulants have been associated with adverse events owing to the non-specific nature of their effects on the coagulation cascade.

Also, there are well-known associations of $\alpha 1$ protease inhibitor deficiency with emphysema and cirrhosis and C1 esterase inhibitor deficiency with angioedema.

We have now found that certain novel amidine compounds are particularly effective as inhibitors of serine proteases, especially proteases with negatively charged P1 specificity pockets, and most especially the serine proteases thrombin, trypsin, urokinase and Factor Xa.

Thus viewed from one aspect we provide serine protease inhibitor compounds having an m-benzamidine group and a lipophilic group coupled to a cyclic group-attached carbon or nitrogen atom (hereinafter the alpha atom), the coupling of the m-benzamidine group to the alpha atom being by a linker group providing a two backbone atom linking chain, the backbone atoms being selected from C, N, O and S and at least one being C, optionally wherein one or both of the backbone atoms form part of a cyclic group and the coupling of the lipophilic group to the alpha atom being by a linker group capable of separating the alpha atom from the

- 3 -

lipophilic group by a range of 2.3 to 6.5 Å in length.

In the compounds of the invention, the lipophilic group is itself optionally substituted by a hydrogen bond donor group. Where this is the case, the lipophilic group and its linker preferably are conformable to separate by from 7.5 to 15.0 Å the alpha atom and the hydrogen bond donor atom of the donor group.


Where distances from the alpha atom to the lipophilic group or to the hydrogen bond donor atom are mentioned, these relate to the distances between the centre of the alpha atom and the centre of the first atom of the lipophilic group or the centre of the hydrogen bond donor atom.

Such distances can be calculated from crystallographic data for any given compound from the bond lengths and bond angles for individual bonds along the length of the molecule between the alpha atom and the first atom of the lipophilic group or the hydrogen bond donor atom. Similarly such distances may be calculated with reasonable accuracy from the bond lengths and bond angles known to be typical of such individual bonds.

The linker between the alpha atom and the lipophilic group may itself be a lipophilic moiety, e.g. an alkylene chain. The nature of the linker may vary considerably - the primary requirement is that it be conformable to place part or all of the lipophilic group at a desired distance away from the alpha atom. The lipophilic group thus desirably is able to occupy at least part of a space 2.3 to 15.0 Å from the alpha atom. The length of the linker generally corresponds to 1 to 5 backbone atoms and may be a chain, branched chain or cyclic linker (e.g. a cyclic amide or an aromatic heterocycle). In one embodiment the alpha atom forms part of a cyclic group which also forms the linker to the lipophilic group.

- 4 -

Thus viewed from an alternative aspect the invention provides serine protease inhibitor compounds of formula I



(where R_1 and R_2 each independently is hydrogen or hydroxyl, alkoxy, alkyl, aminoalkyl, hydroxyalkyl, alkoxyalkyl, alkoxycarbonyl, acyloxymethoxycarbonyl or alkylamino optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl;

15 each R_3 independently is R_1 , R_2 , amino, halo, cyano, nitro, thiol, alkylthio, alkylsulphonyl, alkylsulphenyl, alkylsulphonamido, alkylaminosulphonyl, aminosulphonyl, haloalkoxy and haloalkyl;

20 each X independently is a C, N, O or S atom or a CO, CR_1 , $C(R_1)_2$ or NR_1 group, at least one X being C, CO, CR_1 or $C(R_1)_2$, with the proviso that if the benzamidine group is unsubstituted (i.e. no non-hydrogen R_3 group is present) and the X-X group is $-CH_2C(R_1)_2-$ then R_1 is hydrogen or is attached to the alkylene carbon atom by a heteroatom;

25 L is an organic linker group containing 1 to 5 backbone atoms selected from C, N, O and S, or a branched alkyl or cyclic group;

30 Y (the α -atom) is a nitrogen atom or a CR_1 group or Y and L taken together form a cyclic group;

Cy is a saturated or unsaturated, mono or poly cyclic, homo or heterocyclic group, preferably 35 containing 5 to 10 ring atoms and optionally substituted by groups R_3 or phenyl optionally substituted by R_3 ;

Lp is a lipophilic organic group, e.g. an alkyl,

- 5 -

heterocyclic, alkenyl, alkaryl, cycloalkyl, polycycloalkyl, cycloalkenyl, aryl, aralkyl or haloalkyl group or a combination of two or more such groups optionally substituted by one or more of oxa, oxo, aza, thio, halo, amino, hydroxy or R_3 groups, preferably a group containing up to 25 carbon atoms;

D is a hydrogen bond donor group; and n is 0, 1 or 2);

or a physiologically tolerable salt thereof, e.g. a halide, phosphate or sulphate salt or a salt with ammonium or an organic amine such as ethylamine or meglumine.

In the compounds of the invention, where the alpha atom is carbon it preferably has the conformation that would result from construction from a D- α -aminoacid $NH_2-CR_1(Cy)-COOH$ and a m-carboxyl benzamidine. Likewise the fourth substituent R_1 at an alpha carbon is preferably a methyl or hydroxymethyl group or hydrogen.

In compounds of formula (I) it is envisaged that the unsubstituted or substituted amidino group could be replaced by a substituted or unsubstituted aminomethyl group although an amidino derivative is preferred.

In the compounds of the invention, unless otherwise indicated, aryl groups preferably contain 5 to 10 ring atoms optionally including 1, 2 or 3 heteroatoms selected from O, N and S; alkyl, alkenyl or alkynyl groups or alkylene moieties preferably contain up to 6 carbons; cyclic groups preferably have ring sizes of 3 to 8 atoms; and fused multicyclic groups preferably contain 8 to 16 ring atoms.

The linker group from the benzamidine group to the alpha atom is preferably selected from $-CH=CH-$, $-CONH-$, $-CONR_1-$, $-NH-CO-$, $-NH-CH_2-$, $-CH_2-NH-$, $-CH_2O-$, $-OCH_2-$, $-COO-$, $-OC=O-$ and $-CH_2CH_2-$. Preferably, the X moiety nearest to the alpha atom is an NH or O atom, most preferably a NH group. The X moiety alpha to the phenyl ring is preferably a carbon based group such as CH_2 or


- 6 -

CO, preferably CO. Thus a particularly preferred linker X-X is -CONH-.

The linker group from the alpha atom to the lipophilic group is preferably CO, CH₂NH, CONR₁(CH₂)_m, (CH₂)_mN(R₁)CO(CH₂)_m, (CH₂)_{m+2}, CO(CH₂)_m, (CH₂)_mCO, (CH₂)_mOC=O, (CH₂)_mO or CH=CH(CH₂)_m (where each m is independently 0 or 1). The linker may be optionally branched, for example, to incorporate a polar functionality. In a preferred embodiment Y and L taken together form a cyclic group and the alpha atom is therefore a carbon atom. The cyclic group can be unsubstituted or substituted and can have a ring size of from 3 to 8 atoms. Preferably, the cyclic group is a cyclic amide, most preferably wherein the amide nitrogen of the cyclic amide group is bound to the lipophilic group.

The lipophilic group preferably comprises a cycloalkyl, azacycloalkyl, diazacycloalkyl, phenyl, naphthyl, adamantyl, decalynyl, tetrahydrodecalynyl, bicycloalkyl, mono- or diazabicycloalkyl, mono- or bicyclo heteroaromatic or a linear or branched alkyl, alkylene, alkenyl or alkenylene group all optionally substituted by one or more groups R₃, or a combination of at least two such groups linked by a spiro linkage or a single or double bond or by C=O, O, S, SO, SO₂, CONR₁, NR₁-CO-, NR₁ linkage. For example, representative lipophilic groups include a methyl-cyclohexyl, methylcyclohexylmethyl, methylphenylmethyl, phenylethyl, benzylpiperidinyl, benzoylpiperidinyl, bispiperidinyl or phenylpiperazinyl.

Most preferably, the lipophilic group is selected from



- 8 -

wherein R_3 is R_1 , aryl or cycloalkyl;

m represents 0 or 1;



R_4 represents hydrogen, $(CH_2)_wCOOH$, $(CH_2)_wCONH_2$,
 $(CH_2)_wCON\alpha$ -AminoAcid;

5 w represents an integer from 0 to 4; and


X represents CH or N.

For example specific lipophilic groups include


10




15




20



25



30



35 especially when R_8 represents H, OMe, F, NO_2 , OH or Cl.


- 9 -

The hydrogen bond donor group which may be attached to the lipophilic group preferably has a nitrogen or oxygen atom as the donor atom and conveniently is a hydroxyl group, a primary, secondary or tertiary amine, or a primary or secondary imine group (as part of an amidine or guanidine) or a saturated or unsaturated heterocyclic group containing a ring nitrogen, preferably a group containing 5 to 7 ring atoms. Where the donor atom is a ring nitrogen, the remote portion of the heterocyclic ring may be part of the lipophilic group.

The cyclic group attached to the alpha carbon is preferably an optionally R_3 substituted phenyl or naphthyl group.

The benzamidino group is preferably an unsubstituted m-benzamidino group, or a substituted m-benzamidino group bearing metabolically labile groups such as acyloxymethoxycarbonyl, alkoxycarbonyl or hydroxy.

Accordingly, preferred compounds of the invention are of formula



30

(wherein R_1 , R_2 and R_3 are as hereinbefore defined, R_1 and R_2 preferably being hydrogen or one representing a metabolically labile group such as alkoxycarbonyl or hydroxy, R_3 preferably being hydrogen, OH or NH_2 and when other than hydrogen preferably being para to the amidine group);

35 R_5 and R_6 are hydrogen or taken together with the

- 10 -

carbon atom to which they are attached represent a carbonyl group;

Ar is an unsubstituted or substituted aryl group, preferably phenyl;

5 X-X is -CONH-, -CH₂CH₂-, CH₂O-, -COO-, -CH₂NH-, -OCH₂- or -NHCH₂-;

L₁ is a valence bond or an organic linker group containing 1 to 4 backbone atoms selected from C, N and O;

10 Lp₁ is a cycloalkyl, azacycloalkyl, diazacycloalkyl, phenyl, naphthyl, adamantyl, decaliny, tetrahydrodecaliny, bicycloalkyl, mono- or diazabicycloalkyl, mono- or bicyclo heteroaromatic or a linear or branched alkyl, alkylene, alkenyl or
15 alkenylene group all optionally substituted by a group R₃, or a combination of at least two such groups linked by a spiro linkage or a single or double bond or by C=O, O, S, SO, SO₂, CONR₁, NR₁-CO-, NR₁ linkage. For example, representative lipophilic groups include a methyl-
20 cyclohexyl, methylcyclohexylmethyl, bispiperidinyl, methylphenylmethyl, phenylethyl, benzylpiperidinyl, benzoylpiperidinyl or phenylpiperazinyl and those as hereinbefore described;

D is a hydrogen bond donor group;
25 and n is 0, 1 or 2).

In one embodiment, L₁ comprises the backbone of an alpha amino acid, the lipophilic group being the side chain of the amino acid. The carboxyl part of the alpha amino acid may be optionally coupled via an amide bond
30 to an amino acid or to a primary or secondary cyclic or acyclic alkyl amine or diamine or via an ester bond to primary or secondary alcohols.


In a preferred embodiment, L₁ represents a valence bond and the lipophilic group is bound directly to the
35 carbonyl alpha to the alpha atom via a nitrogen atom which forms part of the lipophilic group. Suitable lipophilic groups in this case therefore include

- 11 -

piperidinyl, pyrrolidinyl and piperazinyl. In a preferred embodiment the piperidine or piperazinyl group is further substituted by a phenyl, benzyl, piperidine, pyridine or benzoyl group, optionally substituted on the phenyl ring by one or more R_3 groups.

In a further embodiment, the lipophilic group has attached a group of the formula $-\text{COOR}_1$ or $-\text{CON-}$ aminoacid or ester derivative thereof.

In another embodiment the group binding the alpha carbon atom to the lipophilic group comprises a heterocyclic group. Accordingly, preferred compounds of the invention also include



20 (wherein R_1 , R_2 and R_3 are as hereinbefore defined R_1 and R_2 preferably being hydrogen or one representing a metabolically labile group such as alkoxycarbonyl or hydroxy, R_3 preferably being hydrogen, OH or NH_2 and when other than hydrogen preferably being para to the amidine group);

25 Ar is an unsubstituted or substituted aryl group, preferably phenyl;

X-X is $-\text{CONH}-$, $-\text{CH}_2\text{CH}_2-$, $\text{CH}_2\text{O}-$, $-\text{COO}-$, $-\text{CH}_2\text{NH}-$, $-\text{OCH}_2-$ or $-\text{NHCH}_2-$;

30 m is 0, 1 or 2;

Het is a 5 or 6-membered heterocyclic group interrupted by 1, 2 or 3 heteroatoms selected from O, N and S optionally substituted by a group R_3 ;

35 Lp_1 is a cycloalkyl, azacycloalkyl, diazacycloalkyl, phenyl, naphthyl, adamantyl, decalinyl, tetrahydrodecalinyl, bicycloalkyl, mono- or diazabicycloalkyl, mono- or bicyclo heteroaromatic or a

- 12 -


linear or branched alkyl, alkylene, alkenyl or
alkenylene group all optionally substituted by a group
 R_3 , or a combination of at least two such groups linked
by a spiro linkage or a single or double bond or by C=O,
5 O, S, SO, SO₂, CONR₁, NR₁-CO-, NR₁ linkage. For example,
representative lipophilic groups include a methyl-
cyclohexyl, methylcyclohexylmethyl, methylphenylmethyl,
phenylethyl, benzylpiperidinyl, benzoylpiperidinyl,
bispiperidinyl or phenylpiperazinyl;

10 D is a hydrogen bond donor group;
and n is 0, 1 or 2).

Where Het is a five membered ring, the two ring
atoms at which it is connected are preferably separated
by one ring atom. Where Het is a six-membered ring, the
15 two ring atoms at which it is connected are preferably
separated by one or two ring atoms. Representative
heterocyclic groups include thiazole, oxazole,
oxadiazole, triazole, thiadiazole or imidazole. Where
the heterocyclic group is substituted by R_3 this is
20 preferably a COOH or COOR₁ connected to the heterocycle
via a valence bond or alkylene chain.

In a further embodiment, the lipophilic group has
attached a group of the formula -COOR₁ or -CON-aminoacid
or ester derivative thereof.

25 The compounds of the invention may be prepared by
conventional chemical synthetic routes, e.g. by amide
bond formation to couple the benzamidine function to the
alpha atom and to couple the lipophilic function to the
alpha atom. Where the alpha atom is a carbon, the
30 cyclic group-alpha atom combination may conveniently
derive from an alpha amino acid with the benzamidine
deriving from a m-amidino-benzoic acid. Amide formation
from such reagents (in which the amidine function may if
desired be protected during some or all of the synthesis
35 steps) yields a compound of formula II.




- 13 -

(where Cy is as defined above and Bd is an optionally protected m-benzamidine group).

The lipophilic group (and optionally simultaneously the hydrogen bond donor) may then conveniently be introduced by reaction of a compound of formula II (or another analogous carboxylic acid) optionally after transformation into an activated form, e.g. an acid chloride or active ester, with a lipophilic group carrying an amine, hydroxylamine, hydrazine or hydroxyl group, e.g. to produce compounds with linkages of -CO-NR₁-, -CO-NR₁-O-, -CO-NR₁-NR₁- and -CO-O- from the alpha atom (where it is a carbon) to the lipophilic group. Where Y and L taken together form a cyclic amide group the lipophilic group can be conveniently introduced by reacting the compound of formula (II) with a lipophilic group carrying a secondary amine with an active side chain. Cyclisation can be base induced via nucleophilic attack of the alpha atom on a leaving group on the active side chain. If necessary the amide linkage can be reduced using an appropriate reducing agent employing the necessary protection depending on whether concurrent reduction of the carboxylic acid moiety is also desired. Alternatively a compound of formula II or another analogous carboxylic acid may be transformed into an alcohol by reaction with isobutylchloroformate and reduction with sodium borohydride.

Such an alcohol, e.g. of formula III



can be reacted to introduce the lipophilic group by reactions such as:

alkylation with an alkyl halide in the presence of a base;

reaction with diethyl azodicarboxylate/triphenylphosphine and a hydroxylated aryl compound;

- 14 -

by reaction with an activated carboxylic acid (e.g. an acid chloride) or with a carboxylic acid and diethylazodicarboxylate/triphenylphosphine;

by reaction with an isocyanate; and

5 by treatment with methanesulphonyl chloride or trifluoromethanesulphonic anhydride and reaction with an amine, or with a thiol optionally followed by oxidation, e.g. with potassium metaperiodate or hydrogen peroxide.

10 In this way compounds with linkages of $-\text{CH}_2-\text{O}-$, $-\text{CH}_2-\text{O}-\text{CO}-$, $-\text{CH}_2-\text{O}-\text{CO}-\text{NR}_1-$, $-\text{CH}_2-\text{NR}_1-$, $-\text{CH}_2-\text{S}-$, $-\text{CH}_2-\text{SO}-$ and $-\text{CH}_2-\text{SO}_2-$ between the alpha carbon and the lipophilic group may be produced.

Alternatively the alcohol can be oxidized to form a corresponding aldehyde (e.g. by oxidation with manganese dioxide or DMSO/oxalyl chloride or DMSO/ SO_3 or Dess-
15 Martin reagent) which may be reacted to introduce the lipophilic group by reactions such as:

reaction with Wittig reagents or Horner-Emmons reagents, optionally followed by reduction of the
20 resulting carbon:carbon double bond using H_2/Pd -carbon;

reaction with an organometallic, eg a Grignard reagent, optionally followed by reaction on the resulting hydroxyl group, such as oxidation (eg with MnO_2 , DMSO/oxalyl chloride or Dess-Martin reagent),
25 alkylation (eg with an alkyl halide in the presence of a base in a solvent such as DMF), arylation (eg with diethylazo dicarboxylate/triphenyl phosphine and a hydroxyaryl compound), ester formation (eg with an acid chloride or with a carboxylic acid and diethylazido
30 dicarboxylate/triphenyl phosphine), or carbamate formation (eg with an isocyanate);

by reaction with an amine followed by reduction, e.g. with sodium cyanoborohydride;

by reaction with a hydrazine; or

35 by reaction with a carbazide.

In this way compounds with linkages of $-\text{CH}=\text{CR}_1-$, $-\text{CH}_2-\text{CHR}_1-$, $-\text{CHOH}-$, $-\text{CHR}_1-\text{O}-$, $-\text{CHR}_1-\text{O}-\text{CO}-$, $-\text{CHR}_1-\text{O}-\text{CO}-\text{NR}_1-$,

- 15 -

-CO-, -CH₂-NR₁-, -CH=N-NR₁- and -CH=N-NR₁-CO-NR₁- between the alpha carbon and the lipophilic group may be produced.

5 The transformation of alcohol to amine referred to above may be used to produce an amine reagent for lipophilic group introduction, e.g. a compound Bd-CONH-CH(Cy)-CH₂-NR₁H.

10 Such an amine reagent may be reacted to introduce the lipophilic group, e.g. by acylation with an acid halide or activated ester, by reaction with isocyanate, by reaction with an isothiocyanate, or by reaction with a sulphonyl chloride. In this way compounds with linkages of -CH₂NR₁-CO-, -CH₂-NR₁-CO-NR₁-, -CH₂NR₁-CS-NR₁- and -CH₂NR₁-SO₂- between the alpha carbon and the
15 lipophilic groups may be produced.

The transformation of acid to amide referred to above may be used to produce an amide reagent for introduction of the lipophilic group, e.g. a compound Bd-CONH-CH(Cy)-CON(R₁)₂.

20 Such amides may be reacted to introduce lipophilic groups, e.g. by reaction with a halo ketone (e.g. phenacyl bromide). This provides a linkage



from alpha carbon to lipophilic group.

Analogously the amide may be transformed to a thioamide by reaction with Lawesson's reagent and then
30 reacted with a halo ketone to form a linkage




35 The amide reagent may likewise be transformed to a nitrile reagent by dehydration, e.g. with trifluoroacetic anhydride. The nitrile reagent may be

- 16 -


reacted with hydrazine then with acyl halide and then cyclized, (e.g. with trifluoroacetic anhydride) to produce a linkage

5



Alternatively it may be treated with hydroxylamine then reacted with acyl halide and cyclized (e.g. with trifluoroacetic anhydride) to produce a linkage


10



The hydrazide produced by reaction of a carboxylic acid reagent with hydrazine discussed above may likewise be used as a reagent for lipophilic group introduction, e.g. as a compound of formula $\text{Bd}-\text{CONH}-\text{CH}(\text{Cy})-\text{CO}-\text{NR}_1-\text{N}(\text{R}_1)_2$.

Thus the hydrazide reagent can be reacted with an acyl halide and cyclized, e.g. with trifluoroacetic anhydride to yield a linkage

20




25

or reacted with an acyl halide or an isocyanate to yield linkages $-\text{CO}-\text{NR}_1-\text{NR}_1-\text{CO}-$ and $-\text{CO}-\text{NR}_1-\text{NR}_1-\text{CO}-\text{NR}_1-$ respectively.

Alternatively the hydrazide may be transformed by reaction with Lawesson's reagent and then reacted with an acyl halide and cyclized (e.g. with trifluoroacetic anhydride) to produce the linkage

35



- 17 -


An alternative route to these compounds is to carry out any of the above chemical reactions to incorporate the lipophilic group (an optional H bond donor) into a protected intermediate such as a compound of formula (IV).



PG = Protecting group

The protecting group may then be removed before coupling of the m-amidino benzoic acid (optionally protected).

A starting reagent for lipophilic group introduction where the alpha atom is nitrogen may be produced for example by reaction of a beta protected hydrazine (such protection to be chosen as to be compatible with the subsequent reagents to be employed) with phosgene, diphosgene, triphosgene or N,N'-carbonyl diimidazole to give a reactive compound of the type:



PG = Protecting group

This intermediate may be used as has been described above for the carboxylic starting reagents where the alpha atom is carbon.


Removal of the protecting group by standard methods and coupling with an activated m-carboxyl-benzamidine will give compounds of the type

- 18 -



(where Bd, X, Y, Cy, L, Lp and D are as defined above).

Thus viewed from a further aspect the invention provides a process for the preparation of a compound according to the invention which process comprises coupling a lipophilic group to a compound of formula (V)



(where Bd, X, Y and Cy are as defined above and Z is a reactive functional group), and optionally subsequently coupling a hydrogen bond donor group to said lipophilic group.

The compounds of the invention may be administered by any convenient route, e.g. into the gastrointestinal tract (e.g. rectally or orally), the nose, lungs, musculature or vasculature or transdermally. The compounds may be administered in any convenient administrative form, e.g. tablets, powders, capsules, solutions, dispersions, suspensions, syrups, sprays, suppositories, gels, emulsions, patches etc. Such compositions may contain components conventional in pharmaceutical preparations, e.g. diluents, carriers, pH modifiers, sweeteners, bulking agents, and further active agents. Preferably the compositions will be sterile and in a solution or suspension form suitable for injection or infusion. Such compositions form a further aspect of the invention. Viewed from this aspect the invention provides a pharmaceutical composition comprising a serine protease inhibitor according to the invention together with at least one pharmaceutically acceptable carrier or excipient.

Viewed from a further aspect the invention provides the use of a serine protease inhibitor according to the invention for the manufacture of a medicament for use in a method of treatment of the human or non-human animal body (e.g. a mammalian, avian or reptilian body) to combat (i.e. treat or prevent) a condition responsive to

- 19 -

said inhibitor.

Viewed from a further aspect the invention provides a method of treatment of the human or non-human animal body (e.g. a mammalian, avian or reptilian body) to combat a condition responsive to a serine protease inhibitor (e.g. a condition such as a thrombotic disorder responsive to a factor Xa inhibitor), said method comprising administering to said body an effective amount of a serine protease inhibitor according to the invention.

The dosage of the inhibitor compound of the invention will depend upon the nature and severity of the condition being treated, the administration route and the size and species of the patient. However in general, quantities of from 0.01 to 100 $\mu\text{mol/kg}$ bodyweight will be administered.

All publications referred to herein are hereby incorporated by reference.

The invention will now be described further with reference to the following non-limiting Examples.

Experimental

Abbreviations used follow IUPAC-IUB nomenclature. Additional abbreviations are Hplc, high-performance liquid chromatography; DMF, dimethylformamide; DCM, dichloromethane; HOAt, 1-hydroxy-7-azabenzotriazole; HATU, [O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate]; Fmoc, 9-Fluorenylmethoxycarbonyl; HOBt, 1-hydroxybenzotriazole; TBTU, 2-(1H-(benzotriazol-1-yl)-1,1,3,3-tetramethyluroniumtetrafluoroborate; DIPEA, diisopropylethylamine; Boc, tertiary butyloxycarbonyl; DIPCI, diisopropylcarbodiimide; DBU, 1,8-diazabicyclo[5.4.0]undec-7-ene; TEA, triethylamine; Rink linker, p-[(R,S)- α -[1-(9H-Fluoren-9-yl)methoxyformamido]-2,4-dimethoxybenzyl]phenyl acetic acid; TFA,

- 20 -

trifluoroacetic acid; MALDI-TOF, Matrix assisted laser desorption ionisation - time of flight mass spectrometry; and RT, retention time. Unless otherwise indicated, amino acid derivatives, resins and coupling reagents were obtained from Novabiochem (Nottingham, UK) and other solvents and reagents from Rathburn (Walkerburn, UK) or Aldrich (Gillingham, UK) and were used without further purification.

Purification: Purification was by gradient reverse phase Hplc on a Waters Deltaprep 4000 at a flow rate of 50 ml/min. using a Deltapak C18 radial compression column (40 mm x 210 mm, 10-15 mm particle size). Eluant A consisted of aqTFA (0.1%) and eluant B 90% MeCN in aqTFA(0.1%) with gradient elution (Gradient 1, 0 min. 20%B then 20% to 100% over 36 min., Gradient 2, 0 min. 5%B for 1 min. then 5%B to 20%B over 4 min., then 20% to 60% over 32 min. or Gradient 3, 0 min. 20%B then 20% to 100% over 15 min.). Fractions were analysed by analytical Hplc and MALDI-TOF before pooling those with >95% purity for lyophilisation.

Analysis: Analytical Hplc was on a Shimadzu LC6 gradient system equipped with an autosampler, a variable wavelength detector at flow rates of 0.4 ml/min. Eluents A and B as for preparative Hplc. Columns used were Techogell5 C18 (2x150mm) (Hplc Technology), Jupiter 5 C18 column (2.1x150 mm, 5µm particle size) (Phenomenex) and Kromasil C4 (2.0x150 mm, 5 mm) (HPLC Technology). Purified products were further analysed by MALDI-TOF and nmr.

All Fmoc-protected amino acids were purchased where available or prepared by known literature methods (1) with the exception of the following novel amino acid:

- 21 -

Preparation of (D,L)-N-Fluorenylmethyloxycarbonyl-4-phenylphenylglycine.

A solution of 4-biphenylcarboxaldehyde (4.6g 25mmol),
5 sodium cyanide (3.68g 75mmol) and ammonium carbonate
(9.60g 100mmol) in 50% aqueous ethanol (175ml) was
heated at 50°C for 20 hours. The reaction mixture was
cooled, concentrated under reduced pressure and
acidified to pH 2.0 with concentrated hydrochloric acid
10 (fume hood). The intermediate 4-(4-phenylphenyl)-2,5-
imidazolidinedione was collected by filtration, washed
with dilute (0.5%) HCl and dried before using as crude
in the next stage. (Filtrates were retained and treated
with sodium hypochlorite solution before disposal)
15 The intermediate 4-(4-phenylphenyl)-2,5-imidazolidinedione
was refluxed in 16% aqueous sodium hydroxide (100ml, 16%
w/v) for 24h. The reaction mixture was then filtered,
cooled, diluted with water (100ml) and then shaken with
ethyl acetate and separated. The aqueous solution was
20 adjusted to pH 5.1 with concentrated hydrochloric acid
and the solid obtained collected by filtration, washed
with a little water and dried to give 4-
phenylphenylglycine (2.36g, 42%). ¹H nmr (d₆ DMSO)
consistent with desired product.
25 To a vigorously stirred solution of 4-
phenylphenylglycine (500mg 2.2mmol) in DCM (20ml) was
added DIPEA (614μl 4.4mmol) and then, carefully,
chlorotrimethylsilane (558μl 4.4mmol) and the mixture
refluxed for 1.5h. The reaction mixture was cooled in
30 an ice bath and fluorenylmethoxycarbonyl chloride (742mg
2.2mmol) was added in one portion. After stirring at 0°C
for 20min the ice bath was removed and stirring was
continued for 1.5h. The reaction mixture was
concentrated under reduced pressure and the residue
35 stirred with a mixture of diethyl ether (20ml) and
saturated sodium carbonate solution (30ml). A yellow
solid which failed to dissolve in either layer was taken

- 22 -

up in water (20ml) and acidified to pH 1 with dilute hydrochloric acid. The mixture was then extracted with ethyl acetate and the organic layer washed with water (2 x 20ml), dried with magnesium sulphate and evaporated to dryness. Recrystallisation from ethanol/water afforded (D,L)-N-fluorenylmethyloxycarbonyl-4-phenylphenylglycine (640mg, 65%).

¹H nmr (d₆ DMSO) 8.25 (1H, d, aromatics); 7.89 (2H, d, aromatics); 7.76 (2H d, aromatics); 7.67 (3H, d, aromatics); 7.57-7.25 (6H, m, aromatics); 5.23 (1H, d, -NH); ~4.27 (3H, m, H_α + CH₂).

Preparation of 3-Amidinobenzoic acid TFA salt

3 Cyanobenzoic acid (10g, 68 mmol) was refluxed in ethanol (300ml) on an isomantle fitted with reflux condenser and soxhlet extractor, the thimble being filled with A4 molecular sieve. Reflux was continued for 10 hours. The heating was then removed and the solution allowed to cool. The solution was then cooled in an ice bath and saturated with hydrogen chloride gas. The sealed flask was allowed to stand overnight then evaporated to dryness. To the dry product was added saturated ammonia/ethanol solution (400ml) and the flask sealed and allowed to stand overnight. The solution was then evaporated to dryness and then treated with 2M sodium hydroxide solution (3 eq., 102ml), the resulting solution was stirred for 2 hours then extracted with ethylacetate (100ml). The aqueous layer was then acidified, with 10% aq hydrochloric acid (200ml) in one lot and extracted with ethylacetate (100ml). Concentrated ammonia solution was added to pH 14 and the solution cooled in the fridge overnight. The precipitate formed was filtered off and washed with water, dissolved in 10%TFA water and lyophilised to a white powder (12g).

- 23 -

¹H nmr (D₂O) 8.40 (1H, s); 8.30 (1H, d, J=7.5Hz); 8.00 (1H, d, J=7.5Hz); 7.72 (1H, t).

Synthesis of inhibitors

5

Method 1: Using a solid phase strategy on a Protein Technologies, Symphony Multiple Peptide Synthesiser by attachment of bis amino compounds to Peg-2-chlorotrityl chloride resin: 2-Chlorotrityl chloride resin was typically treated with greater than 2 fold excess of the di-amine in dry DCM. The resin was further modified by the attachment of acids. Activation of Fmoc protected amino acid (2-5eq) was by TBTU/ DIPEA, all couplings (minimum 120 min.) were carried out in DMF. Deprotection of the Fmoc group was achieved with 20% (v/v) piperidine in DMF. Other acid substituents were added as the HOBt or HOAt esters either by activation with HBTU/HATU or DIPCI with or without Boc protection of amino groups. Cleavage of the products from the resin was by treatment (30 min., ambient) with 10% (v/v) triethylsilane in TFA, filtration, evaporation and trituration with diethylether.

Synthesis using the Symphony Multiple Peptide Synthesiser.

The Symphony Multiple Peptide Synthesiser is charged with DMF, DCM, TBTU in DMF (450 mM), DIPEA in DMF (900 mM), 20% (v/v) piperidine in DMF. Resins are held in plastic reaction vessels that allow the introduction of reagents and solvents and nitrogen for agitation or air drying.

A typical synthesis cycle on the Symphony is as follows:-

- 24 -

The reaction vessel containing the resin (0.1 mmol) is charged with the Fmoc protected amino acid (0.5 mmol) and then this is dissolved in DMF (2.5ml), treated with TBTU (0.56 mmol, 1.25ml) and DIPEA (1.1 mmol, 1.25ml) and agitated with nitrogen for 2 hours (agitation times may vary). After coupling the resin is washed with DMF (6x 5ml) then deprotected with 20% (v/v) piperidine in DMF (2x 5ml for 1 min.each, then 1x 5ml for 8 min.) the resin is then washed with DMF (6x 5ml).

Example 1:**3-Amidino-benzoyl-D-phenylglycine 4-aminomethylcyclohexyl methylamide**

Bis-1,4 aminomethylcyclohexane(2ml) was added to 2 chlorotrityl chloride resin (1.2 mmol/g, 0.73g) pre swollen in dry DCM (4ml). After 2h the resin was washed with DCM (6x5ml), DMF (6x5ml) and DCM (6x5ml). The resin was then air dried to allow aliquots to be taken.

The bis-1,4-aminomethylcyclohexane 2-chlorotrityl chloride resin (0.1 mmol) was treated with Fmoc-D-Phenylglycine (0.5 mmol, 187mg), DMF(2.5ml), TBTU in DMF(1.25ml of a 450mM solution) and DIPEA in DMF (1.25ml of a 900 mM solution). The mixture was agitated with nitrogen for 2 hours. Deprotection and washing as above.

HOBt (0.5 mmol, 68 mg) dissolved in DMF(4ml) at 0°C was treated with DIPCI (0.5 mmol, 80µl), for 10 min.

3-amidinobenzoic acid TFA salt (0.5 mmol, 139 mg.) was added and stirring continued for 10 min. at room temperature. The mixture was transferred to the reaction vessel on the Symphony and agitated for 10 hours with nitrogen. The resin was washed with DMF (6x5ml), DCM (6x5ml) and air dried. The product was cleaved from the resin with 10% triethylsilane in TFA (10ml) for 30 minutes, the resin filtered off and the TFA solution evaporated to dryness and triturated with diethyl ether

- 25 -

to give the crude product. The crude product was dissolved in water (10ml), filtered and purified by preparative reverse phase Hplc.

5 ¹H nmr (D₂O) Mixture of cyclohexyl cis and trans isomers
8.09 (1H, s); 8.05 (1H, d, J=7.5Hz); 7.90 (1H, d,
J=7.5Hz); 7.66 (1H, t, J=7.5Hz); 7.43 (5H, m); 5.47 (1H,
s); 3.05 (2H, m); 2.78 (2H, m); 1.48 (7H, m); 0.86 (3H,
m) MS TOF 422 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1,
10 water/acetonitrile/TFA) rt 13.45 min. (major) and 13.62
min. (minor).

Compounds made by the above method:-

15 **Example 2.**

3-Amidino-benzoyl-DL-3-chloro-phenylglycine-4-aminomethylcyclohexyl methylamide

¹H nmr (D₂O) Mixture of cyclohexyl cis and trans isomers
8.19 (1H, s); 8.15 (1H, d, J=7.5Hz); 8.04 (1H, d,
20 J=7.5Hz); 7.90 (1H, t, J=7.5Hz); 7.53 (4H, m); 5.63 (1H,
s); 3.18 (2H, m); 2.90 (2H, m); 1.66 (7H, m); 1.00 (3H,
m). (MS TOF 457 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2,
water/acetonitrile/TFA) rt 16.53 min.

25 **Example 3.**

3-Amidino-benzoyl-DL-4-methoxyphenylglycine-4-aminomethylcyclohexyl methylamide

¹H nmr (D₂O) Mixture of cyclohexyl cis and trans isomers
8.19 (1H, s); 8.12 (1H, d, J=7.5Hz); 8.01 (1H, d, J-
30 7.5Hz); 7.80 (1H, t, J=7.5Hz); 7.50 (2H, d); 7.10,
(2H, d); 5.53 (1H, s); 3.86 (3H, s); 3.13 (2H, m); 2.90
(2H, m); 1.66 (7H, m); 0.95 (3H, m). (MS TOF 452 (M+1⁺).
Hplc (Jupiter5 C18, Gradient 2, water/acetonitrile/TFA)
rt 13.87 min. (major) and 14.10 min. (minor).

35

- 26 -

Example 4.**3-Amidino-benzoyl-DL-4-hydroxyphenylglycine 4-aminomethylcyclohexyl methylamide**

¹H nmr (D₂O) Mixture of cyclohexyl cis and trans isomers
5 7.40 (1H, d); 7.25 (2H, dd); 7.08 (1H, d); 7.02 (1H, d); 6.88 (2H, m); 6.82 (1H, d); 5.28 (1H, d); 5.10 (1H, d); 3.03 (2H, m); 2.83 (2H, m); 1.50 (7H, m); 0.85 (3H, m). MS TOF 438 (M+1⁺) Hplc (Jupiter5 C18, Gradient 2, water/acetonitrile/TFA) rt 9.69 min. (major) and 14.10
10 min. (minor).

Example 5.**3-Amidino-benzoyl-DL-1-naphthylglycine 4-aminomethylcyclohexyl methylamide**

¹H nmr (D₂O) Mixture of cyclohexyl cis and trans
15 isomers 8.40 (1H, s); 8.25 (4H, t); 8.132 (1H, d, J=7.5Hz); 7.81 (4H, m); 6.53 (1H, s); 3.70-3.10 (2H, m); 2.99 (2H, m); 2.0-1.3 (7H, m); 1.07 (3H, m). (MS TOF 472 (M+1⁺)). Hplc (Jupiter5 C18, Gradient 2, water/acetonitrile/TFA) rt 17.62 min. (major) and 18.03
20 min. (minor).

Example 6.**3-Amidino-benzoyl-DL-4-phenyl phenylglycine 4-aminomethylcyclohexyl methylamide**

¹H nmr (D₂O) Mixture of cyclohexyl cis and trans isomers
25 8.11 (2H, m); 7.94 (1H, d, J=7.5Hz); 7.71 (5H, m); 7.51 (5H, m); 5.58 (1H, s); 3.3-2.90 (2H, m); 2.90-2.65 (2H, m); 1.8-1.2 (7H, m); 0.85 (3H, m). MS TOF 498 (M+1⁺)
30 Hplc (Jupiter5 C18, Gradient 2, water/acetonitrile/ TFA) rt 19.37 min. (major), 21.69 min. (minor).

Example 7.**3-Amidino-benzoyl-DL-4-trifluoromethyl phenylglycine 4-aminomethylcyclohexyl methylamide**

¹H nmr (D₂O) Mixture of cyclohexyl cis and trans isomers
35 8.12 (1H, s); 8.07 (1H, d, J=7.5Hz); 7.93 (1H, d,

- 27 -

J=7.5Hz); 7.8-7.6 (5H, m); 5.61 (1H, s); 3.30-2.70 (4H, m); 1.50 (7H, m); 0.85 (3H, m). MS TOF 490 (M+1⁺) Hplc (Jupiter5 C18, Gradient 2, water/acetonitrile/TFA) rt 19.37 min.

5

Example 8.**3-Amidino-benzoyl-R,S-3-phenyl β -alanine 4-aminomethylcyclohexyl methylamide**

¹H nmr D₂O Mixture of cyclohexyl cis and trans isomers.

10 8.08 (2H, m); 7.96 (1H, d, J=7.5Hz); 7.73 (1H, t, J=7.5Hz); 7.44 (5H, m); 5.43 (1H, t, J=7.9Hz); 3.50 (2H, m); 2.88 (4H, m); 1.6-1.0 (7H, m); 0.77 (2H, m). MS TOF 436 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2, water/acetonitrile/TFA) rt 13.22 min. (major), 13.61
15 (minor)

Example 9.**3-Amidino-benzoyl-DL-3-indolyglycine 4-aminomethylcyclohexyl methylamide**

20 ¹H nmr (D₂O) Mixture of cyclohexyl cis and trans isomers
8.1-7.85 (3H, m); 7.65 (1H, m); 7.47 (4H, m); 7.33 (1H, m); 4.85 (m, partially obscured by solvent); 4.3-3.80 (4H, m); 3.05 (2H, m); 2.78 (3H, m); 1.8-1.0 (7H, m); 1.0-0.65 (3H, m). MS TOF 451 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2, water/acetonitrile/TFA) rt 10.12 min.
25

Example 10.**3-Amidino-benzoyl-DL-piperonylglycine 4-aminomethylcyclohexyl methylamide**

30 ¹H nmr D₂O Mixture of cyclohexyl cis and trans isomers.
8.13 (2H, m); 8.00 (1H, d, J=7.5Hz); 7.72 (1H, t, J=7.5Hz); 6.96 (5H, m); 5.98 (2H, s); 5.44 (1H, s); 3.4-2.7 (4H, m); 1.9-1.25 (7H, m); 1.1-0.8 (3H, m). MS TOF 466 (M+1⁺) Hplc (Kromasil C4, Gradient 2, water/acetonitrile/ TFA) rt 12.62 min.
35

- 28 -

Example 11.**3-Amidino-benzoyl-DL-3-methyl phenylglycine 4-aminomethylcyclohexyl methylamide**¹H nmr D₂O Mixture of cyclohexyl cis and trans isomers.

5 8.09 (2H, m); 7.96 (1H, d, J=7.5Hz); 7.70 (1H, t, J=7.5Hz); 7.25 (4H, m); 5.50 (1H, s); 3.4-2.7 (4H, m); 2.31 (3H, s); 1.9-1.1 (7H, m); 1.1-0.7 (3H, m). MS TOF 436 (M+1⁺) Hplc (Kromasil C4, Gradient 2, water/acetonitrile/ TFA) rt 14.41 min.

10

Example 12.**3-Amidino-benzoyl-DL-2-naphthylglycine 4-aminomethylcyclohexyl methylamide**¹H nmr D₂O Mixture of cyclohexyl cis and trans isomers.

15 7.98 (2H, m); 7.83 (4H, m); 7.56 (1H, t, J=7.5Hz); 7.45 (4H, m); 5.58 (1H, s); 3.25-2.55 (4H, m); 1.7-0.9 (7H, m); 0.9-0.5 (3H, m). MS TOF 472 (M+1⁺) Hplc (Kromasil C4, Gradient 2, water/acetonitrile/ TFA) rt 17.44 min. (major) 17.69 (minor).

20

Example 13.**3-Amidino-benzoyl-1-aminocyclopentane-N- 4-aminomethylcyclohexyl methyl-1-carboxamide**¹H nmr D₂O Mixture of cyclohexyl cis and trans isomers.

25 7.92 (2H, m); 7.81 (1H, d, J=7.5Hz); 7.57 (1H, t, J=7.5Hz); 3.6-3.1 (2H, m); 2.91 (2H, d, J=6.8Hz); 2.68 (2H, d, J=6.8Hz); 2.2-1.8 (4H, m); 1.8-1.2 (13H, m); 1.0-0.65 (3H, m). MS TOF 400 (M+1⁺) Hplc (Kromasil C4, Gradient 2, water/acetonitrile/ TFA) rt 10.87 min.

30

Example 14.**3-Amidino-benzoyl-DL-cyclohexylglycine 4-aminomethylcyclohexyl methylamide**¹H nmr (D₂O) Mixture of cyclohexyl cis and trans isomers

35 8.10 (1H, s); 8.05 (1H, d, J=7.5Hz); 7.85 (1H, d, J=7.5Hz); 7.66 (1H, t, J=7.5Hz); 4.25 (1H, d); 3.05 (2H, m); 2.78 (2H, m); 1.48 -0.86 (19H, m) MS TOF 428 (M+1⁺).

- 29 -

Hplc (Jupiter5 C18, Gradient 2, water/acetonitrile/TFA)
rt 14.76 min. (major) and 15.09 min. (minor).

Example 15.

5 **3-Amidino-benzoyl-L-phenylglycine 4-
aminomethylcyclohexyl methylamide**
MS TOF 422 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2,
water/acetonitrile/ TFA) rt 10.06 min.

10 **Example 16.**

**3-Amidino-benzoyl-D-phenylglycine 3-
aminomethylcyclohexyl methylamide**
MS TOF 422 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2,
water/acetonitrile/ TFA) rt 8.81 min.

15

Example 17.

**3-Amidino-benzoyl-L-phenylglycine 3-
aminomethylcyclohexyl methylamide**
MS TOF 422 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2,
20 water/acetonitrile/ TFA) rt 8.79 min.

Example 18.

**3-Amidino-benzoyl-DL-2-thienylglycine 4-
aminomethylcyclohexyl methylamide**
25 MS TOF 429 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2,
water/acetonitrile/TFA) rt 12.73 min.

Example 19.

**3-Amidino-benzoyl-DL-4-chloro-phenylglycine 4-
aminomethylcyclohexyl methylamide**
30 MS TOF 457 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2,
water/acetonitrile/TFA) rt 15.89 min.

Example 20

35 **3-Amidino-benzoyl-D-phenylglycine 4-
aminomethylbenzylamide**
MS TOF 416 (M+1⁺) Hplc (Techogel15 C18, Gradient 2,

- 30 -

water/acetonitrile/ TFA) rt 12.34 min.

Example 21.

3-Amidino-benzoyl-D-phenylglycine 3-aminomethylbenzylamide

MS TOF 416 (M+1⁺) Hplc (Techogel15 C18, Gradient 2, water/acetonitrile/ TFA) rt 9.65 min.

Example 22.

3-Amidino-benzoyl-L-phenylglycine 3-aminomethylbenzylamide

MS TOF 416 (M+1⁺) Hplc (Techogel15 C18, Gradient 2, water/acetonitrile/ TFA) rt 10.26 min.

Example 23.

3-Amidino-benzoyl-L-phenylglycine 4-aminomethylbenzylamide

¹H nmr (D₂O) 8.10 (2H, m); 7.88 (1H, d, J=7.5Hz); 7.65 (1H, t, J=7.5Hz); 7.4-7.1 (9H, m); 5.57 (1H, s); 4.37 (2H, ABq); 4.06 (2H, s). MS TOF 416 (M+1⁺) Hplc (Techogel15 C18, Gradient 1, water/acetonitrile /TFA) rt 12.34 min.

Example 24.

3-Amidino-benzoyl-D-phenylglycine -4-(4-aminocyclohexyl methyl) cyclohexylamide

¹H nmr (D₂O) 8.3-7.95 (3H, m); 7.85-7.6 (2H, m); 7.47 (4H, m); 5.55 (1H, s); 3.59 (1H, m); 3.06 (1H, m); 2.1-1.5 (9H, m); 1.5-0.85 (11H, m). MS TOF 490 (M+1⁺). Hplc (Kromasil C4, Gradient 2, water/acetonitrile /TFA) rt 18.30 min.

Example 25.

3-Amidino-benzoyl-L-phenylglycine -4-(4-aminocyclohexyl methyl) cyclohexylamide

¹H nmr (D₂O) 8.3-7.95 (3H, m); 7.85-7.6 (2H, m); 7.47 (4H, m); 5.55 (1H, s); 3.59 (1H, m); 3.06 (1H, m); 2.1-

- 31 -

1.5 (9H, m); 1.5-0.85 (11H, m). MS TOF 490 (M+1⁺). Hplc (Kromasil C4, Gradient 2, water/acetonitrile /TFA) rt 18.39 min.

5 **Example 26.**

3-Amidino-benzoyl-D-phenylglyciny-D-2-naphthylalanine-5-aminopentamide

¹H nmr (d₄methanol) d 8.21 (1H, s); 8.14 (1H, d, J=7.5Hz); 7.96 (1H, d, J=7.5Hz); 7.80-7.65 (6H, m);
10 7.45-7.15 (7H, m); 5.66 (1H, s); 4.56 (1H, m); 3.16 (2H, m); 3.02 (2H, m); 2.68 (2H, m); 1.45 (2H, m); 1.32 (2H, m); 1.12 (2H, m). MS TOF 579 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA) rt 13.56 min.

15 **Example 27.**

3-Amidino-benzoyl-D-phenylglyciny D-lysine-5-aminopentamide

¹H nmr (D₂O) d 8.10 (1H, s); 8.05 (1H, d, J=7.5Hz); 7.94 (1H, d, J=7.5Hz); 7.68 (1H, t, J=7.5Hz); 7.45 (5H, m);
20 5.53 (1H, s); 4.22 (1H, t); 3.05 (2H, m); 2.90 (4H, dd); 1.8-1.0 (12H, m). MS TOF 510 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA) rt 10.56 min.

Example 28.

25 **3-Amidino-benzoyl-D-phenylglycine-5-aminopentamide**

¹H nmr (D₂O) d 7.85 (1H, s); 7.80 (1H, d, J=7.5Hz); 7.70 (1H, d, J=7.5Hz); 7.55 (1H, t, J=7.5Hz); 7.20 (5H, m);
5.30 (1H, s); 3.05 (2H, m); 2.60 (2H, m); 1.25 (4H, m); 0.90 (2H, m); MS TOF 382 (M+1⁺). Hplc (Jupiter5 C18,
30 Gradient 2, water/acetonitrile/TFA) rt 11.26 min.

Example 29.

3-Amidino-benzoyl-D-phenylglycine-R,S-3-aminopyrrolidinamide

35 ¹H nmr (D₂O) 8.18 (1H, s); 8.10 (1H, d); 7.88 (1H, d); 7.70 (1H, t); 7.50 (5H, m); 5.55 (1H, s); 4.50 (1H, m); 3 - 4 (4H, m); 2.33 (1H, m); 2.00 (1H, m). MS TOF

- 32 -

366(M+1⁺). Hplc (Jupiter5 C18, Gradient 2, water/acetonitrile/TFA) rt 10.38 + 10.79 min.

Example 30.

5 **3-Amidino-benzoyl-D-phenylglycine 4-aminomethyl piperidinamide**

MS TOF 394 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2, water/acetonitrile/TFA) rt 11.40 min. (major) and 11.97 min. (minor).

10

Method 2: By solid phase strategy on a Protein Technologies, Symphony Multiple Peptide Synthesiser using Fmoc amino acids attached to Peg-2-chlorotrityl chloride resin: Typically the 2-chlorotrityl chloride resin was treated with a 2 fold excess of the Fmoc amino acid in a 1:1 mixture of DMF and dry DCM and DIPEA (2eq.). The resin was washed with DMF/DCM and deprotected with 20% piperidine in DMF before further modification. The resin was further modified by the attachment of acids. Activation of Fmoc protected amino acids (2-5eq) was by activation with TBTU/ DIPEA, all couplings (minimum 120 min.) were carried out in DMF. Deprotection of the Fmoc group was achieved with 20% (v/v) piperidine in DMF. Other acid substituents were added as the HOBt or HOAt esters either by activation with HBTU/HATU or DIPCI with or without Boc protection of amino groups. Cleavage of the products from the resin was by treatment (30 min., ambient) with 10% (v/v) triethylsilane in TFA, filtration, evaporation and trituration with ether.

30

Example 31.

3-Amidino-benzoyl-D-phenylglyciny-D-2-naphthylalaninyl-glycine

35

Fmoc Glycine (0.2mmol, 59 mg.) in DMF (2ml) was added to 2 chlorotrityl chloride resin (1.0 mmol/g, 0.1g) pre

- 33 -

swollen in dry DCM (2ml), then DIPEA (0.2 mmol). After 2h the resin was washed with DCM (6x5ml), DMF (6x5ml) and DCM (6x5ml). The resin was then air dried to allow aliquots to be taken for further modification.

5

On the Symphony Fmoc-Glycyl-2-chlorotrityl resin (0.1 mmol) was deprotected with 20% (v/v) piperidine in DMF and washed with DMF(6x 5ml) then treated with Fmoc-D-2-naphthylalanine (0.5 mmol 220 mg), DMF(2.5ml), TBTU in 10 DMF(1.25ml of a 450mM solution) and DIPEA in DMF (1.25ml of a 900 mM solution). The mixture was agitated with nitrogen for 2 hours. Deprotection and washing as above. The resin was then treated Fmoc-D-Phenylglycine (0.5 mmol 187 mg), DMF(2.5ml), TBTU in DMF(1.25ml of a 450mM 15 solution) and DIPEA in DMF (1.25ml of a 900 mM solution). The mixture was agitated with nitrogen for 2 hours. Deprotection and washing as above.

HOBt (0.5 mmol, 68 mg) dissolved in DMF(4ml) was stirred at 0°C with DIPCI (0.5 mmol, 80µl), for 10 min. 20 3-amidinobenzoic acid TFA salt (0.5 mmol, 139 mg.) was added and stirring continued for 10 min. at room temperature. The mixture was transferred to the reaction vessel on the Symphony and agitated for 10 hours with nitrogen. The resin was then washed with DMF (6x5ml), 25 DCM (6x5ml) and air dried. The product was cleaved from the resin with 10% (v/v) triethylsilane in TFA (10ml) for 30 minutes, the resin filtered off, the TFA solution evaporated to dryness and triturated with diethyl ether to give the crude product. The crude product was then 30 dissolved in water (10ml), filtered and purified by preparative reverse phase Hplc..

¹H nmr (DMSO) δ 8.15 (1H, s); 8.12 (1H, d, J=7.5Hz); 7.85 (1H, d, J=7.5Hz); 7.74 (5H, m) ;7.65 (1H, t); 7.42 - 35 7.25 (7H,m); 5.73 (1H, s); 4.72 (1H, m); 3.75(2H, d); signals at 3.3-3.0 obscured by solvent. MS TOF 552 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA)

- 34 -

rt 14.88 min.

Compounds made by the above method:-

5 **Example 32.**

**3-Amidino-benzoyl-D-phenylglycinyl-D-2-naphthylalaninyl
N-methylglycine**

¹H nmr (CD₃CN) mixture of rotomers, major product quoted.
d 7.95 (2H, m); 7.87 (1H, d, J=7.5Hz); 7.71 (4H, m);
10 7.62 (1H, t, J=7.5Hz); 7.36 (3H, m); 7.22 (5H, m); 5.57
 (1H, s); 5.19 (1H, dd); 3.95 (2H, ABq); 2.98 (3H, s).
MS TOF 566 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1,
water/acetonitrile/TFA) rt 14.86 min.

15 **Example 33.**

**3-Amidino-benzoyl-D-phenylglycinyl-D-2-naphthylalaninyl-
D-proline**

¹H nmr (CD₃CN) d 8.15 (1H, s); 8.08 (1H, d); 7.95 (1H,
d); 7.82 (5H, m); 7.62 , (1H,m); 7.22 - 7.5 (7H,m); 5.63
20 (1H, s); 5.05 (1H, m); 4.35(1H, m); 3.60 , (2H,m);
 signals at 3.3 obscured by solvent; 3.1- , (1H,m); 2.22,
 (1H,m); 2.0 , (4H,m); . MS TOF 592 (M+1⁺). Hplc (Jupiter5
C18, Gradient 1, water/acetonitrile/TFA) rt 15.32 min.

25 **Method 3:** By solid phase strategy on a Protein

Technologies, Symphony Multiple Peptide Synthesiser
using Fmoc amino acids attached to TentaGel S- resin
(Rapp Polymere) via the Rink amide linker: Typically the
TentaGel resin was treated with a 5 fold excess of the
30 Rink linker, TBTU (1eq.), DIPEA (2eq.). The resin was
 washed with DMF and deprotected with 20% piperidine in
 DMF before further modification. The resin was further
 modified by the attachment of acids. Activation of Fmoc
 protected amino acid (2-5eq) was by TBTU/ DIPEA, all
35 couplings (minimum 120 min.) were carried out in DMF.
 Deprotection of the Fmoc group was achieved with 20%
 piperidine in DMF. Other acid substituents were added as

- 35 -

the HOBt or HOAt esters either by activation with HBTU/HATU or DIPCI with or without Boc protection of amino groups. Cleavage of the products from the resin was by treatment (30 min., ambient) with 10% triethylsilane in TFA, filtration, evaporation and trituration with ether.

Example 34.**3-Amidino-benzoyl-D-phenylglyciny l D-phenylalanine amide**

10

On the Symphony TentaGel S -NH₂ resin (0.1 mmol, 400 mg, 0.24 mmol./g) was treated with Rink linker (0.5 mmol, 270mg), DMF(2.5ml), TBTU in DMF(1.25ml of a 450mM solution) and DIPEA in DMF (1.25ml of a 900 mM solution). The mixture was agitated with nitrogen for 2 hours. Deprotection and washing as above.

15

On the Symphony Rink-TentaGel resin (0.1 mmol) was then treated with Fmoc-D-Phenylalanine- (0.5 mmol, 194mg), DMF(2.5ml), TBTU in DMF(1.25ml of a 450mM solution) and DIPEA in DMF (1.25ml of a 900 mM solution). The mixture was agitated with nitrogen for 2 hours. Deprotection and washing as above.

20

The resin (0.1 mmol) was then treated with Fmoc-D-Phenylglycine (0.5 mmol, 187mg), DMF(2.5ml), TBTU in DMF(1.25ml of a 450mM solution) and DIPEA in DMF (1.25ml of a 900 mM solution). The mixture was agitated with nitrogen for 2 hours. Deprotection and washing as above.

25

HOBt (0.5 mmol, 68 mg) dissolved in DMF(4ml) was stirred at ice bath with DIPCI (0.5 mmol, 80μl) for 10 min. 3-amidino-benzoic acid TFA salt (0.5 mmol, 139 mg.) was added and stirring continued for 10 min. at room temperature. The mixture was then transferred to the reaction vessel on the Symphony and agitated for 10 hours with nitrogen. The resin was then washed with DMF (6x5ml), DCM (6x5ml) and air dried. The product was

35

- 36 -

cleaved from the resin with 10% triethylsilane in TFA (10ml) for 30 minutes, the resin filtered off, the TFA solution evaporated to dryness and triturated with diethyl ether to give the crude product. The crude product was then dissolved in water (10ml), filtered and purified by preparative reverse phase Hplc.

¹H nmr (d₄methanol) δ 8.18 (1H, s); 8.10 (1H, d, J=7.5Hz); 7.87 (1H, d, J=7.5Hz); 7.72 (1H, t, J=7.5Hz); 7.40 (5H, m); 7.10 (5H, m); 5.53 (1H, s); 4.55 (dd, partially obscured by solvent); 3.08 (1H, dd); 2.87 (1H, dd). MS TOF 444 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA) rt 11.70 min.

Compounds made by the above method:-

Example 35.

3-Amidino-benzoyl-D-phenylglycinyll-L-asparagine amide

¹H nmr (D2O) δ 8.10 (1H, s); 8.04 (1H, d, J=7.5Hz); 7.91 (1H, d, J=7.5Hz); 7.66 (1H, t, J=7.5Hz); 7.42 (5H, m); 5.47 (1H, s); 4.73 (m, partially obscured by solvent); 2.71 (2H, m); 2.59 (2H, m). MS TOF 411 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2, water/acetonitrile/TFA) rt 10.49 min.

Example 36.

3-Amidino-benzoyl-D-phenylglycinyll-D-2-naphthylalanine amide

¹H nmr (d₄methanol) δ 8.18 (1H, s); 8.12 (1H, d, J=7.5Hz); 7.95 (1H, d, J=7.5Hz); 7.74 (5H, m); 7.42 (3H, m); 7.25 (5H, m); 5.63 (1H, s); 3.12 (1H, m); signals at ~4.8 and 3.3 obscured by solvent. MS TOF 494 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA) rt 15.46 min.

- 37 -

Example 37.**3-Amidino-benzoyl-D-phenylglycinyll-D-valine amide**

¹H nmr (d₄methanol) d 8.44 (1H, s); 8.36 (1H, d, J=7.5Hz); 8.10 (1H, d, J=7.5Hz); 7.86 (1H, t, J=7.5Hz); 7.70 (2H, m); 7.55 (3H, m); 5.96 (1H, s); 4.40 (1H d); 2.21 (1H, m); 1.13 (6H, t).). MS TOF 396 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA) rt 5.71 min.

Example 38.**3-Amidino-benzoyl-D-phenylglycinyll-D-lysine amide**

¹H nmr (D₂O) d 8.10 (1H, s); 8.04 (1H, d, J=7.5Hz); 7.95 (1H, d, J=7.5Hz); 7.66 (1H, t, J=7.5Hz); 7.43 (5H, m); 5.56 (1H, s); 4.26 (1H, m); 2.91 (2H, t); 1.77 (2H, m); 1.62 (2H, m); 1.42 (2H, m). MS TOF 425 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2, water/acetonitrile/TFA) rt 10.73 min.

Example 39.**3-Amidino-benzoyl-D-phenylglycinyll-L-lysine amide**

¹H nmr (D₂O) d 8.08 (1H, s); 8.00 (1H, d, J=7.5Hz); 7.99 (1H, d, J=7.5Hz); 7.61 (1H, t, J=7.5Hz); 7.40 (5H, m); 5.52 (1H, s); 4.21 (1H, m); 2.68 (2H, m); 1.74 (1H, m); 1.60 (1H, m); 1.40 (2H, m); 1.08 (1H, m). MS TOF 425 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2, water/acetonitrile/TFA) rt 10.61 min.

Example 40.**3-Amidino-benzoyl-D-phenylglycinyll-D,L-phenylglycinyll-L-valine amide**

¹H nmr (CD₃CN) mixture of diastereomers d 8.39 (1H, m); 8.15 (1H, m); 7.93 (1H, m); 7.70 (1H, m); 7.41 (10H, m); 5.71 (1H, d); 5.46 (1H, m); 4.14 (1H, m); 2.77 (1H, m); 0.97 (3H, m); 0.65 (3H, m). MS TOF 529 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA) rt 13.10 min.

- 38 -

Example 41.**3-Amidino-benzoyl-D-phenylglycinyll-D-asparagine-D-asparagine amide**

¹H nmr (D₂O) d 8.28 (1H, s); 8.20 (1H, d, J=7.5Hz); 8.08
5 (1H, d, J=7.5Hz); 7.82 (1H, t, J=7.5Hz); 7.59 (5H, m);
5.72 (1H, s); 4.70 (1H, dd); 2.85 (4H, m).. MS TOF 526
(M+1⁺). Hplc (Jupiter5 C18, Gradient 2,
water/acetonitrile/TFA) rt 11.14 min.

Example 42.**3-Amidino-benzoyl-D-phenylglycinyll-D-leucine amide**

¹H nmr (CD₃CN) d 8.23 (1H, s); 8.14 (1H, d, J=7.5Hz);
7.93 (1H, d, J=7.5Hz); 7.72 (1H, t, J=7.5Hz); 7.48 (5H,
m); 5.62 (1H, s); 4.32 (1H, m); 1.58 (2H, m); 0.87 (6H,
15 dd). MS TOF 410 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2,
water/acetonitrile/TFA) rt 14.98 min.

Example 43.**3-Amidino-benzoyl-D-phenylglycinyll-L-phenylalanine amide**

¹H nmr (CD₃CN) d 8.20 (1H, s); 8.14 (1H, d, J=7.5Hz);
7.96 (1H, d, J=7.5Hz); 7.72 (1H, t, J=7.5Hz); 7.35 (5H,
m); 7.14 (3H, m); 7.04 (2H, m); 5.60 (1H, s); 5.46 (1H,
dd); 3.15 (1H, dd); 2.80 (1H, m). MS TOF 444 (M+1⁺). Hplc
(Jupiter5 C18, Gradient 1, water/acetonitrile/TFA) rt
25 12.78 min.

Example 44.**3-Amidino-benzoyl-D-phenylglycinyll-L-2-naphthylalanine amide**

¹H nmr (d₄methanol) d 8.26 (1H, s); 8.19 (1H, d,
J=7.5Hz); 7.95 (1H, d, J=7.5Hz); 7.80 (1H, m); 7.67 (3H,
m); 7.53 (1H, s); 7.45 (2H, m); 7.25 (1H, d); 7.08 (3H,
m); 6.92 (2H, t); 5.61 (1H, s); 3.45 (1H, m); 3.0 (1H, m)
signal at ~4.8 obscured by solvent. MS TOF 494 (M+1⁺).
35 Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA)
rt 16.17 min.

- 39 -

Example 45.**3-Amidino-benzoyl-D-phenylglyciny-L-valine amide**

¹H nmr (CD₃CN) d 8.17 (1H, s); 8.08 (1H, d, J=7.5Hz);
7.89 (1H, d, J=7.5Hz); 7.65 (1H, t, J=7.5Hz); 7.37 (5H,
5 m); 5.64 (1H, s); 4.08 (1H d); 1.97 (m, partially
obscured by solvent); 0.63 (6H, t). MS TOF 396 (M+1⁺).
Hplc (Jupiter5 C18, Gradient 2, water/acetonitrile/TFA)
rt 9.34 min.

10 **Example 46.****3-Amidino-benzoyl-D-phenylglyciny-D-asparagine amide**

¹H nmr (D₂O) d 7.91 (1H, s); 7.86 (1H, d, J=7.5Hz); 7.63
(1H, d, J=7.5Hz); 7.48 (1H, t, J=7.5Hz); 7.23 (5H, m);
5.30 (1H, s); 4.42 (m, partially obscured by solvent);
15 2.57 (2H, m). MS TOF 411 (M+1⁺). Hplc (Jupiter5 C18,
Gradient 2, water/acetonitrile/TFA) rt 10.23 min.

Example 47.20 **3-Amidino-benzoyl-L-phenylglyciny-D-2-naphthylalanine
amide**

¹H nmr (CD₃CN/D₂O) 8.17 (1H, s); 8.10 (1H, d); 7.92 (1H,
d); 7.81 (1H, m); 7.67 (3H, m); 7.53 (1H, s); 7.48 (2H,
m); 7.22 (1H, d); 7.12 (3H, m); 6.98 (2H, m); 5.56
(1H,s); 4.70 (1H, dd); 3.35 (1H, dd); 3.00 (1H,dd). MS
25 TOF 494 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1,
water/acetonitrile/TFA) rt 15.46 min.

Example 48.30 **3-Amidino-benzoyl-L-phenylglyciny-L-2-naphthylalanine
amide**

¹H nmr (CD₃CN) 8.07 (1H, s); 8.01 (1H, d); 7.92 (1H, d);
7.78 (1H, m); 7.66 (2H, s); 7.41 (4H, m); 7.14 (6H, d);
5.46 (1H,s); 4.73 (1H, dd); 3.38 (1H, dd); 3.08 (1H,dd).
MS TOF 494 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1,
35 water/acetonitrile/TFA) rt 14.51 min.

- 40 -

Example 49.**3-Amidino-benzoyl-D-phenylglycinyll-D-4-chloro
phenylalanine amide**

¹H nmr (CD₃CN) δ 8.39 (1H, s); 8.32 (1H, d, J=7.5Hz);
5 8.13 (1H, d, J=7.5Hz); 7.90 (1H, t, J=7.5Hz); 7.51 (5H,
m); 7.40 (4H, ABq); 5.70 (1H, s); 4.77 (1H, dd); 3.36
(1H, dd); 3.09 (1H, dd). MS TOF 479 (M+1⁺). Hplc
(Jupiter5 C18, Gradient 2, water/acetonitrile/TFA) rt
13.63 min.

10

Example 50.

3-Amidino-benzoyl-D-phenylglycinyll-D-isoleucine amide
MS TOF 410 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2,
water/acetonitrile/TFA) rt 14.36 min.

15

Example 51.

3-Amidino-benzoyl-D-phenylglycinyll-D-tyrosine amide
MS TOF 460 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2,
water/acetonitrile/TFA) rt 12.56 min.

20

Example 52.

**3-Amidino-benzoyl-D-phenylglycinyll-D-1-naphthylalanine
amide**
MS TOF 494 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2,
25 water/acetonitrile/TFA) rt 22.19 min.

Example 53.

3-Amidino-benzoyl-D-phenylglycinyll-D-threonine amide
MS TOF 398 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2,
30 water/acetonitrile/TFA) rt 10.63 min.

Example 54.

3-Amidino-benzoyl-D-phenylglycinyll-D-histidine amide
MS TOF 434 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2,
35 water/acetonitrile/TFA) rt 10.04 min.

- 41 -

Example 55.

3-Amidino-benzoyl-D-phenylglyciny-D-phenylalanine amide
MS 444 TOF ($M+1^+$). Hplc (Jupiter5 C18, Gradient 2,
water/acetonitrile/TFA) rt 14.82 min.

5

Example 56.

3-Amidino-benzoyl-D-phenylglyciny-D-phenylalaniny D-proline amide
MS TOF 541 ($M+1^+$). Hplc (Jupiter5 C18, Gradient 2,
water/acetonitrile/TFA) rt 11.54 min.

10

Example 57.

3-Amidino-benzoyl-D-phenylglyciny-D-tryptophan amide
MS TOF 483 ($M+1^+$). Hplc (Jupiter5 C18, Gradient 2,
water/acetonitrile/TFA) rt 17.14 min.

15

Method 4: By solution phase strategy: Typically an activated Boc-amino acid was treated with an amine (primary or secondary) or alcohol (1eq.). Activation of Boc protected amino acid was by HATU or TBTU/DIPEA(1:2), all couplings (minimum 120 min.) were carried out in DMF. After an aqueous work up the deprotection of the Boc group was achieved with TFA. Other acid substituents were added as the HOBt or HOAt esters either by activation with HBTU/HATU or DIPCI with or without Boc protection of amino groups. The final products were purified by preparative reverse phase Hplc.

20

25

30

Example 58.

3-Amidino-benzoyl-D-phenylglycine -1-adamantylamide

Boc D-phenylglycine (251 mg, 1 mmol.) was dissolved in DMF(3ml) with HATU (380 mg., 1 mmol.) and DIPEA(350 μ l ., 2 mmol.). To this mixture was added 1-adamantylamine hydrochloride(187 mg., 1 mmol.) and DIPEA (170 μ l., 1 mmol.). The mixture was stirred overnight. The mixture

35

- 42 -

was then taken up into ethylacetate and washed with water, sodium carbonate solution, water, 10% hydrochloric acid solution and water. The ethylacetate was evaporated without drying and treated immediately with TFA for 30 min. The TFA was then evaporated to dryness and the product triturated with diethylether. TEA(1ml) was added and evaporated to dryness. HOBt (1 mmol, 136 mg) dissolved in DMF(4ml) was stirred at 0°C and treated with DIPCI (1 mmol, 160 μ l), for 10 min. 3 amidino benzoic acid TFA salt (1 mmol., 278 mg.) was added and stirring continued for 10 min. at room temperature. The mixture was then added to the D-phenylglycine adamantyl amide and stirred overnight. The crude product was dissolved in water/acetonitrile (20ml), filtered and purified by preparative Hplc to yield pure product (150 mg.)

¹H nmr (D₂O/CD₃CN) 8.21 (1H, s); 8.13 (1H, d, J=7.5Hz); 7.92 (1H, d, J=7.5Hz); 7.70 (1H, t, J=7.5Hz); 7.46 (2H, m); 7.34 (3H, m); 5.52 (1H, s); 2.15-1.52 (15H, m). M.S. TOF 431 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/ TFA) rt 19.54 min.

Compounds made by the above method:-

Example 59.

3-Amidino-benzoyl-D-phenylglycine -1-adamantylmethanamide

¹H nmr (D₂O/CD₃CN) 8.23 (1H, s); 8.15 (1H, d, J=7.5Hz); 7.85 (1H, d, J=7.5Hz); 7.71 (1H, t, J=7.5Hz); 7.46 (2H, m); 7.35 (3H, m); 5.59 (1H, d); 3.0 (2H, s obscured by solvent); 2.15-1.52 (15H, m). M.S. TOF 445 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/ TFA) rt 21.35 min.

- 43 -

Example 60.**3-Amidino-benzoyl-D-phenylglycine-2-phenylethylamide**

¹H nmr (d₄ methanol) 8.18 (2H, m); 7.87 (1H, d, J=7.5Hz);
7.60 (1H, t, J=7.5Hz); 7.30 (5H, m); 7.02 (5H, m); 5.53
5 (1H, s); 3.32 (1H, m); 1H signal at ~3.25 obscured by
solvent); 2.68 (2H, t). M.S. TOF 401 (M+1⁺). Hplc
(Kromasil C4, Gradient 1, water/acetonitrile/TFA) rt
15.56 min.

10 **Example 61.****3-Amidino-benzoyl-D-phenylglycine-2-(1-adamantyl) ethyl ester**

¹H nmr (d₄ methanol) 8.27 (2H, m); 7.94 (1H, d J=7.5Hz);
7.72 (1H, t, J=7.5Hz); 7.42 (5H, m); 5.66 (1H, s); 4.25
15 (2H, m); 1.82 (3H, s); 1.61 (6H, m); 1.38 (8H, m). M.S.
TOF 459 (M+1⁺). Hplc (Kromasil C4, Gradient 1,
water/acetonitrile/ TFA) rt 23.17 min.

Example 62.20 **3-Amidino-benzoyl-D-phenylglycine-2-methylbenzylamide**

¹H nmr (CD₃CN) 8.14 (2H, m); 7.88 (1H, d, J=7.5Hz); 7.65
(1H, t, J=7.5Hz); 7.45 (2H., m); 7.35 (3H, m); 7.04 (4H,
m); 5.59 (1H, s); 4.30 (2H, AB q); 2.13 (3H, s). M.S.
TOF 401 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1,
25 water/acetonitrile/ TFA) rt 14.81 min.

Example 63.**3-Amidino-benzoyl-D-phenylglycine-4-benzoyl piperidinamide**

30 ¹H nmr (DMSO) 8.40 (2H, m); 8.10 (1H, d); 7.70 (1H, t);
7.50 (10H, m); 5.55 (1H, s); 3.60 (1H, m); 2.5 (2H,
m); 1.00 (6H, m); .S. TOF 469 (M+1⁺). Hplc (Jupiter5 C18,
Gradient 1, water/acetonitrile/ TFA) rt 11.43 min.

35 **Example 64.****3-Amidino-benzoyl-D-phenylglycine-4-methylbenzylamide**

¹H nmr (CD₃CN) 8.30 (1H, s); 8.22 (1H, d, J=7.5Hz); 7.95

- 44 -

(1H, d, J=7.5Hz); 7.71(1H, t, J=7.5Hz); 7.51 (2H., m); 7.34 (3H, m); 7.04 (4H, s); 5.76 (1H, s); 4.30 (2H, AB q); 2.30 (3H, s). M.S. TOF 401 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/ TFA) rt 15.20 min.

5

Example 65.**3-Amidino-benzoyl-D-phenylglycine-1-naphthylmethylamide**

¹H nmr (CD₃CN/D₂O) 8.14 (1H, s); 8.02 (1H, d, J=7.5Hz); 7.84 (3H, m); 7.60 (1H, d, J=7.5Hz); 7.32 (5H, m, J=7.5Hz); 7.12 (5H., m); 5.58 (1H, s); 4.50 (2H, AB q); 3.03 (2H, s). M.S. TOF 436 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/ TFA) rt 16.77 min.

10

Example 66.**3-Amidino-benzoyl-D-phenylglyciny-D-phenylalanine -N-pyrrolidinamide**

¹H nmr (CD₃CN/D₂O) 8.14 (1H, s); 8.03 (1H, d, J=7.5Hz); 7.94 (1H, d, J=7.5Hz); 7.70(1H, t, J=7.5Hz); 7.42 (5H, m); 7.42 (5H., m); 5.62 (1H, s); 4.80 (1H, t); 3.40 , (1H, m); 3.20 (2H, m); 3.02 , (2H, d); 2.80 , (1H, m); 1.65 , (4H, m); . M.S. TOF 499 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/ TFA) rt 14.03 min.

20

Example 67.**3-Amidino-benzoyl-D-phenylglycine-3-methylbenzylamide**

¹H nmr (CD₃CN) 7.95 (1H, s); 7.81 (1H, d, J=7.5Hz); 7.60 (1H, d, J=7.5Hz); 7.45 (1H, t, J=7.5Hz); 7.23 (2H., m); 7.12 (3H, m); 6.60-6.75 (4H, m); 5.40 (1H, s); 4.10 (2H, AB q); 1.95 (3H, s). M.S. TOF 401 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/ TFA) rt 14.52 min.

30

Example 68.**3-Amidino-benzoyl-D-phenylglycine-2adamantylamide**

MS TOF 431 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 12.68 min.

35

Method 5: By solid phase strategy using Boc amino acids

- 45 -

attached to Kaiser Oxime resin (Novabiochem): Typically the resin is treated with a 2 fold excess of the Boc phenylglycine symmetrical anhydride. The resin was washed with DCM/DMF then DCM and deprotected with 25%TFA in DCM before further modification. The resin was further modified by the attachment of acids. Activation of Boc protected amino acid (2-5eq) was by TBTU/ DIPEA, all couplings (minimum 120 min.) were carried out in DCM/DMF. Deprotection of the Boc group was achieved with 25%TFA in DCM. Other acid substituents were added as the HOBt or HOAt esters either by activation with HBTU/HATU or DIPCI with or without Boc protection of amino groups. Cleavage of the products from the resin was by treatment (1 day, ambient) with an amine or alcohol in chloroform, washing with chloroform and methanol/chloroform.

Example 69.**3-Amidino-benzoyl-D-phenylglycine cyclohexyl amide**

Boc-D-Phenylglycine (0.5 mmol, 126 mg) was dissolved in DCM (5ml) and treated with DIPCI (0.5 mmol., 80 μ l) and stirred for 10 min. A white precipitate forms which was dissolved by adding DMF. This mixture was added to Oxime resin (0.11 mmol, 330 mg, 0.33 mmol./g) previously swelled with DCM. 4-Dimethyl amino pyridine (10 mg, cat.) was added and the mixture swirled for 2 hours. The resin was filtered and washed with DMF/DCM then DCM. The resin was then treated with 25% TFA/DCM (20ml) for 20 min., filtered and washed with DCM. HOBt (0.5 mmol, 68 mg) dissolved in DMF(4ml) was stirred at 0°C with DIPCI (0.5 mmol, 80 μ l) for 10 min. 3 amidino-benzoic acid TFA salt (0.5 mmol, 139 mg.) was added and stirring continued for 10 min. at room temperature. The mixture was then added to the phenylglycine-oxime resin and agitated for 2 hours. The resin was filtered washed with DMF, DCM and chloroform

- 46 -

then treated with cyclohexylamine (0.3 mmol, 40 μ l) in chloroform (2ml) and agitated for 1 day. The resin was then filtered off and washed with chloroform (10ml) and chloroform/methanol (1:1, 10ml). The combined organic
5 extracts were evaporated to dryness and washed with 10% aqueous acetic acid (5-10ml). The aqueous insoluble residue was dissolved in acetonitrile/water and lyophilised.

MS TOF 379 ($M+1^+$). Hplc (Jupiter5 C18, Gradient 3,
10 water/acetonitrile/TFA) rt 10.62 min.

Example 70.

3-Amidino-benzoyl-D-phenylglycine-3,3-dimethylbutylamide

MS TOF 382 ($M+1^+$). Hplc (Jupiter5 C18, Gradient 3,
15 water/acetonitrile/TFA) rt 13.90 min.

Example 71.

3-Amidino-benzoyl-D-phenylglycine-1,2,3,4-tetrahydro-1-naphthylamide

MS TOF 427 ($M+1^+$). Hplc (Jupiter5 C18, Gradient 3,
20 water/acetonitrile/TFA) rt 11.74 min.

Example 72.

3-Amidino-benzoyl-D-phenylglycine-R,S 3-methyl-2-butylamide

MS TOF 367 ($M+1^+$). Hplc (Jupiter5 C18, Gradient 3,
25 water/acetonitrile/TFA) rt 10.21 + 10.42 min.

Example 73.

3-Amidino-benzoyl-D-phenylglycine-3-phenylpropylamide

MS TOF 415 ($M+1^+$). Hplc (Jupiter5 C18, Gradient 3,
30 water/acetonitrile/TFA) rt 14.00 min.

Example 74.

3-Amidino-benzoyl-D-phenylglycine-3-trifluoromethylbenzylamide

MS TOF 455 ($M+1^+$). Hplc (Jupiter5 C18, Gradient 3,

- 47 -

water/acetonitrile/TFA) rt 14.32 min.

Example 75.

3-Amidino-benzoyl-D-phenylglycine-3-fluorobenzylamide

5 MS TOF 405 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 12.56 min.

Example 76.

3-Amidino-benzoyl-D-phenylglycine-2-methyl propylamide

10 MS TOF 353 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 11.10 min.

Example 77.

3-Amidino-benzoyl-D-phenylglycine-4-phenyl butylamide

15 MS TOF 429 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 14.96 min.

Example 78.

3-Amidino-benzoyl-D-phenylglycine-cyclohexylmethylethylamide

20 MS TOF 493 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 11.52 min.

Example 79.

3-Amidino-benzoyl-D-phenylglycine-2,2 diphenylethylamide

25 MS TOF 478 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 12.65 min.

Example 80.

3-Amidino-benzoyl-D-phenylglycine-R-1-(1-naphthyl)ethylamide

30 MS TOF 450 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 12.26 min.

Example 81.

3-Amidino-benzoyl-D-phenylglycine-S-1-(2-naphthyl)ethylamide

35 MS TOF 450 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,

- 48 -

water/acetonitrile/TFA) rt 11.19 min.

Method 6: By solution phase strategy: Typically an activated Fmoc amino acid was treated with an alcohol (1eq.) which contains a Boc protected amino group. Activation of Fmoc protected amino acid was by HATU or TBTU/ DIPEA(1:2), all couplings (minimum 120 min.) were carried out in DMF with DBU as catalyst After an aqueous work up the deprotection of the Fmoc group was achieved with 20% piperidine in DMF followed by evaporation and column chromatography. Other acid substituents were added as the HOBT or HOAt esters either by activation with HBTU/HATU or DIPCI with or without Boc protection of amino groups. The products were purified by preparative reverse phase Hplc, followed by deprotection of the amino groups with TFA. Further modification to acetamidine or amidine can be achieved by known literature methods. (2,3)

Example 82.

3-Amidino-benzoyl-D-phenylglycine -(4-piperidinyl)methyl ester

Fmoc D-Phenylglycine(1 mmol., 374 mg) was dissolved in DMF(5ml) with HATU (1 mmol., 380 mg) and DIPEA(2 mmol., 350 μ l). To this mixture was added N-Boc 4-hydroxymethyl piperidine (215 mg, 1 mmol.) and DBU (100 μ l). The mixture was stirred overnight. The mixture was then taken up into ethylacetate(50ml) and washed with water, sodium carbonate solution, water, 10% hydrochloric acid solution and water. The ethylacetate layer was dried over magnesium sulphate, filtered and evaporated. The product was then treated with 20% piperidine in DMF for 90 min. this was then evaporated to dryness and then dissolved in methanol and adsorbed onto silica gel 60. The material was then chromatographed on silica gel 60, eluting first with hexane/ethylacetate (2:1) then

- 49 -

hexane/ethylacetate (1:2), the required product eluting in the second solvent. The combined fractions were evaporated to dryness.

5 HOBt (1 mmol, 136 mg) dissolved in DMF(4ml) was stirred at 0°C with DIPCI (1 mmol, 160 μ l) for 10 min. 3 amidino-benzoic acid TFA salt (1 mmol., 278 mg.) was added and stirring continued for 10 min. at room temperature. The mixture was then added to the D-phenylglycine N-Boc-4-piperidine methyl ester and 10 stirred overnight. The mixture was evaporated to dryness on high vacuum, water and acetonitrile added to allow full dissolution and the whole lyophilised. The resulting crude product was treated with TFA/water (9:1) for 30 min. and evaporated to dryness. The crude product. 15 was dissolved in water/acetonitrile (20ml), filtered and purified by preparative reverse phase Hplc. Producing pure product (200mg).

¹H nmr (D₂O) 8.12 , (1H,m); 8.05 (1H, d); 7.90 (1H, d); 20 7.65 (1H, t); 7.45 (5H, m); 5.65 , (1H,s); 4.10 , (2H,d); 3.30 (2H,m); 2.80 (2H, m); 1.90 (1H, m); 1.75 (2H, m); 1.30 (2H, m); MS TOF 395 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2, water/acetonitrile/TFA) rt 12.48 min.

25 Compounds made by the above method:-

Example 83

3-Amidino-benzoyl-D-phenylglycine-2-(4-piperidinyl) ethyl ester

30 ¹H nmr (D₂O) 8.12 , (1H,m); 8.05 (1H, d); 7.90 (1H, d); 7.65 (1H, t); 7.45 (5H, m); 5.65 , (1H,s); 4.25 , (2H,m); 3.20 (2H,m); 2.60 (2H, m); 1.65 (2H, m); 1.50 (2H, m); 1.20 (3H, m); MS TOF 409 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2, water/acetonitrile/TFA) rt 13.14 min.

35

- 50 -

Example 84.**3-Amidino-benzoyl-D-phenylglycine-2-(N-acetimino-4-piperidinyl) ethyl ester**

¹H nmr (D₂O) 8.22 , (1H,m); 8.20 (1H, d); 8.02 (1H, d);
5 7.80 (1H, t); 7.60 (5H, m); 5.75 , (1H,s); 4.35 , (2H,m);
3.85 (2H,m); 3.00 (3H, m); 2.30 (3H,s); 1.65 (3H, m);
1.50 (1H, m); 1.20 (2H, m); MS TOF 450 (M+1⁺). Hplc
(Jupiter5 C18, Gradient 2, water/acetonitrile/TFA) rt
14.92 min.

10

Example 85.**3-Amidino-benzoyl-D-phenylglycine-(N-acetimino-4-piperidinyl) methyl ester**

¹H nmr (D₂O) 8.12 , (1H,m); 8.05 (1H, d); 7.90 (1H, d);
15 7.65 (1H, t); 7.40 (5H, m); 5.60 , (1H,s); 4.05 , (2H,m);
3.75 (2H,m); 3.10 (1H, m); 2.10 (3H,d); 1.90 (1H, m);
1.60 (2H, m); 1.20 (2H, m); 0.90 (1H,m); MS TOF 436
(M+1⁺). Hplc (Jupiter5 C18, Gradient 2,
water/acetonitrile/TFA) rt 13.34 min.

20

Example 86.**3-Amidino-benzoyl-D-phenylglycine-(N-amidino-4-piperidinyl) methyl ester**

¹H nmr (D₂O) 8.02 , (1H,m); 7.97 (1H, d); 7.80 (1H, d);
25 7.55 (1H, t); 7.40 (5H, m); 5.55 , (1H,s); 4.05 , (2H,m);
3.65 (2H,m); 2.80 (2H, m); 1.80 (1H,m); 1.50 (2H, m);
1.10 (1H, m); 0.80 (1H,m); MS TOF 437 (M+1⁺). Hplc
(Jupiter5 C18, Gradient 2, water/acetonitrile/TFA) rt
13.61 min.

30

Example 87.**3-Amidino-benzoyl-D-phenylglycine-2-(N-amidino-4-piperidinyl) ethyl ester**

¹H nmr (D₂O) 8.17 , (1H,m); 8.07 (1H, d); 7.93 (1H, d);
35 7.70 (1H, t); 7.45 (5H, m); 5.60 , (1H,s); 4.25 , (2H,m);
3.55 (2H,m); 2.75 (2H, m); 1.60 (4H,m); 1.25 (1H, m);
1.00 (2H, m); MS TOF 451 (M+1⁺). Hplc (Jupiter5 C18,

- 51 -

Gradient 2, water/acetonitrile/TFA) rt 14.97 min.

Example 88.

2-Hydroxy-5-amidino-benzoyl D-phenylglycine-4-aminomethylcyclohexyl methylamide

5 5-Bromosalicylic acid (9.31mmol., 2.02g) was dissolved in DMF (4.5ml) and treated with copper (I) cyanide (11.33 mmol., 1.2eq., 1.015g) and heated under reflux for 4 hours. The solution was allowed to cool and poured
10 into a solution of ferric chloride (2g) and concentrated hydrochloric acid (0.7ml) in water (3ml). The mixture was heated for 10 min. at 60°C, then cooled and filtered, the solid was washed with water, then recrystallised ethanol/water. Product 0.73g, 48%.

15 On the Symphony bis-1,4-aminomethylcyclohexane 2 chlorotriptyl chloride resin (0.1 mmol) (See method 1) was treated with Fmoc-D-Phenylglycine (0.5 mmol, 187mg), DMF(2.5ml), TBTU in DMF(1.25ml of a 450mM solution) and
20 DIPEA in DMF (1.25ml of a 900 mM solution) and agitated with nitrogen for 2 hours. Deprotection and washing as above.

HOBt (0.5 mmol, 68 mg) dissolved in DMF(4ml) was stirred
25 at 0°C with DIPCI (0.5 mmol, 80µl) for 10 min. 2-hydroxy-5-cyanobenzoic acid (0.5 mmol, 84 mg.) was added and stirring continued for 10 min. at room temperature. The mixture was then transferred to the reaction vessel on the Symphony and agitated for 10 hours with nitrogen.
30 The resin was washed with DMF (6x5ml), DCM (6x5ml) and air dried. The product was cleaved from the resin with 10% triethylsilane in TFA (10ml) for 30 minutes, the resin filtered off, the TFA solution evaporated to dryness and triturated with diethyl ether to give the
35 crude product. The crude product was dissolved in saturated hydrogen chloride in ethanol (10ml), sealed and allowed to stand overnight. The solution was

- 52 -

evaporated to dryness and treated with saturated ammonia in ethanol (10ml) and allowed to stand overnight. The solution was then evaporated to dryness, dissolved in water and purified by reverse phase Hplc.

5

¹H nmr (D₂O), 8.35 , (1H,s); 7.90 , (1H,d); 7.55 , (5H,m); 7.25 , (1H,d); 5.65 (1H, s); 3.20 (2H, m); 2.90 (2H, m); 1.40-1.80 (7H, m); 0.95 (3H, m) MS TOF 438 (M+1⁺). Hplc (Techogel5 C18, Gradient 2, water/acetonitrile/TFA) rt 14.11 min.

10

Example 89.**2-Amino-5-amidinobenzoyl-D-phenylglycine-4-aminomethylcyclohexyl methylamide**

15

2-Amino-5-bromo benzoic acid (1.6 mmol., 345mg) was dissolved in N-methylpyrrolidone (5ml) and treated with copper (I) cyanide (2.39 mmol., 1.5eq., 207mg) under reflux for 4.5 hours. The solution was allowed to cool and poured into water (10ml). The mixture was acidified to pH3 with hydrochloric acid and extracted with ethylacetate, the ethylacetate was washed with water, dried over magnesium sulphate and evaporated to a yellow solid (145mg, 56%).

20

25

On the Symphony bis-1,4-aminomethylcyclohexane-2-chlorotrityl resin (0.1 mmol) (See method 1) was treated with Fmoc-D-Phenylglycine (0.5 mmol, 187 mg), DMF (2.5ml), TBTU in DMF (1.25ml of a 450mM solution) and DIPEA in DMF (1.25ml of a 900 mM solution) for 2 hours. Deprotection and washing as above.

30

HOBt (0.5 mmol, 68 mg) dissolved in DMF (4ml) was stirred at 0°C with DIPCI (0.5 mmol, 80μl) for 10 min. 2-amino-5-cyanobenzoic acid (0.5 mmol, 84 mg.) was added and stirring continued for 10 min. at room temperature. The mixture was transferred to the reaction vessel on the Symphony and agitated for 10 hours with nitrogen. The

35

- 53 -

resin was then washed with DMF (6x5ml), DCM (6x5ml) and air dried. The product was cleaved from the resin with 10% triethylsilane in TFA (10ml) for 30 minutes, the resin filtered off, the TFA solution evaporated to dryness and triturated with diethyl ether to give the crude product. The crude product was then dissolved in saturated hydrogen chloride in ethanol (10ml), sealed and allowed to stand overnight. The solution was then evaporated to dryness and treated with saturated ammonia in ethanol (10ml) and allowed to stand overnight. The solution was then evaporated to dryness, dissolved in water and purified by reverse phase Hplc.

¹H nmr (D₂O) Mixture of cyclohexyl cis and trans isomers 7.80 (1H, s); 7.50 (1H, d, J=7.5Hz); 7.35 (5H, m); 6.75 (1H, d, J=7.5Hz); 5.35 (1H, s); 2.87 (2H, m); 2.60 (2H, m); 1.00-1.60 (7H, m); 0.70 (3H, m) MS TOF 437 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2, water/acetonitrile/TFA) rt 13.11 min. (major), 13.31 (minor).

Prepared by Method 2

Example 90.

3-Amidino-benzoyl-D-phenylglyciny-D-2-naphthylalanine

¹H nmr (CD₃CN) 8.00 (1H, s); 7.95 (1H, d); 7.87 (1H, d); 7.72 (4H, m); 7.63 (2H, m); 7.38 (3H, m); 7.25 (4H, m); 5.60 (1H, s); 4.70 (1H, m); 3.25 (2H, m). MS TOF 495 (M+1⁺). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 10.19 min.

Example 91.

3-Amidino-benzoyl-D-phenylglycine-1,2,3,4, tetrahydroisoquinolin-2-yl-3-carboxylic acid

¹H nmr (DMSO) mixture of isomers and rotomers (only one isomer cited in nmr) 8.28 (1H, s); 8.18 (1H, d); 7.90 (1H, d); 7.69 (1H, t); 7.35 (4H, m); 7.15 (5H, m); 6.30 (1H, s); 5.15, (1H, m); 5.70, (2H, m); 3.15, (2H, d). MS

- 54 -

TOF 457 ($M+1^+$). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA) rt 13.84 and 14.33 min.

Example 92.

5 **3-Amidino-benzoyl-D-phenylglyciny1-D-2-naphthylalaniny1-D-proline**

^1H nmr (CD_3CN) d 8.15 (1H, s); 8.08 (1H, d); 7.95 (1H, d); 7.82 (5H, m); 7.62 (1H, m); 7.22 - 7.5 (7H, m); 5.68 (1H, s); 5.01 (1H, m); 4.25 (1H, m); 3.50 (1H, m);
10 3.1, (2H, m); 2.22 (1H, m); 2.0 (3H, m); . MS TOF 592 ($M+1^+$). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA) rt 16.04 min.

Example 93.

15 **3-Amidino-benzoyl-D-phenylglycine-D-2-naphthylalaniny1-L-pipecolic acid**

^1H nmr (CD_3CN) 7.95 (1H, s); 7.90 (1H, d); 7.85 (1H, d); 7.62 (5H, m); 7.25 (8H, m); 5.60 (1H, s); 5.15 (1H, t); 4.90 (1H, m); 3.60 (1H, m); 3.00 (4H, m); 2.05 (1H, m);
20 1.45 (2H, m); 1.15 (1H, m); 0.08 (1H, m). MS TOF 606 ($M+1^+$). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 14.90 min.

Example 94.

25 **3-Amidino-benzoyl-D-phenylglycine-D-2-naphthylalaniny1-D-pipecolic acid**

^1H nmr (CD_3CN) 7.95 (1H, s); 7.90 (1H, d); 7.85 (1H, d); 7.62 (5H, m); 7.25 (8H, m); 5.45 (1H, s); 5.10 (1H, t); 5.00 (1H, m); obscured by solvent 3.60 (1H, m); 3.00 (4H, m);
30 2.05 (1H, m); 1.45 (2H, m); 1.25 (1H, m); 1.10 (1H, m). MS TOF 606 ($M+1^+$). Hplc (Magellan C8, Gradient 3, water/acetonitrile/TFA) rt 14.71 min.

Example 95.

35 **3-Amidino-benzoyl-D-naphthylglyciny1-D-2-naphthylalaniny1-D-proline**

^1H nmr (CD_3CN) d 8.13 (1H, s); 8.0 (2H, m); 7.85 (8H, m);

- 55 -

7.62 , (1H,t); 7.45 (4H, m); 7.05 (2H,m); 6.35 (1H, s);
5.10 (1H, m); 4.34 (1H, m); 3.67 , (2H,m); 3.20, (2H,m);
2.22 , (1H,m); 2.0 (obscured by solvent) , (3H,m). MS TOF
642 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1,
5 water/acetonitrile/TFA) rt 18.14 min.

Prepared by method 3

Example 96.

10 **3-Amidino-benzoyl-D-naphthylglyciny-D-2-naphthylalanine
amide**

¹H nmr (CD₃CN) d 8.25 (1H, d); 8.13 (1H, s); 7.95 (H, m);
7.85 (H, m); 7.74 , (H,m); 7.65 (1H,t); 7.50 (H,m); 7.39
(2H,d); 6.83 (H,m); 6.22 (1H, s); 4.85 (1H, m);
15 3.25, (2H,m). MS TOF 545 (M+1⁺). Hplc (Jupiter5 C18,
Gradient 1, water/acetonitrile/TFA) rt 17.32 min.

Prepared by method 4

20 **Example 97.**

**3-Amidino-benzoyl-D-phenylglycine-2-(4-
methoxyphenyl)ethylamide**

¹H nmr (D4 methanol) 8.20 (2H, m); 8.00 (1H, d);
7.73 (1H, t); 7.40 (5H, m); 6.85 (4H, m); 5.60 (1H, m);
25 3.75 (2H, m); 2.70 (2H, m); 1.85 (3H, s). MS TOF 431
(M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 10.09 min.

Example 98.

30 **3-Amidino-benzoyl-D-phenylglycine-2-(4-
chlorophenyl)ethylamide**

¹H nmr (D4 methanol) 8.28 (1H, s); 8.21 , (1H,d); 7.97
(1H, d); 7.73 (1H, t); 7.40 (5H, m); 7.10 (4H, m); 5.61
(1H, m); 4.45 (2H, m); 2.78 (2H, m). MS TOF 436 (M+1⁺).
35 Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA)
rt 11.26 min.

- 56 -

Example 99.**3-Amidino-benzoyl-D-phenylglycine-4-chloro benzylamide**

¹H nmr (D4 methanol) 8.20 (1H, s); 8.11 , (1H,d); 7.90 (1H, d); 7.65 (1H, t); 7.50 (5H, m); 7.17 (4H, m); 5.60 (1H, m); 4.31 (2H, m). MS TOF 422 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 10.81 min.

Example 100.**3-Amidino-benzoyl-D-phenylglycine-4-methoxy benzylamide**

¹H nmr (D4 methanol) 8.22 (1H, s); 8.15 , (1H,d); 7.93 (1H, d); 7.68 (1H, t); 7.40 (5H, m); 6.95 (4H, m); 5.63 (1H, m); 4.30 (2H, m); 3.75 , (3H,s). MS TOF 417 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 9.73 min.

Example 101.**3-Amidino-benzoyl-D-phenylglycine-3-chloro benzylamide**

¹H nmr (D4 methanol) 8.32 (1H, s); 8.25 , (1H,d); 7.95 (1H, d); 7.75 (1H, t); 7.50 (5H, m); 7.20 (4H, m); 5.72 (1H, m); 4.40 (2H, m). MS TOF 422 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 10.78 min.

Example 102.**3-Amidino-benzoyl-D-phenylglycine-4,4'-dimethyl dibenzylamide**

¹H nmr (DMSO) 8.30 (1H, s); 8.21 , (1H,d); 7.95 (1H, d); 7.70 (1H, t); 7.45 (5H, m); 7.10 (10H, m); 6.15 (1H, m); 4.41 (4H, m); 2.30 , (6H,d). MS TOF 505 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 14.39 min.

Example 103.**3-Amidino-benzoyl-D-phenylglycine-N-methyl-2-phenethylamide**

¹H nmr (DMSO) 8.25 (1H, s); 8.20 , (1H,d); 7.95 (1H, d); 7.70 (1H, t); 7.30 (10H, m); 6.02 (1H, m); 3.78 (1H, m); 3.54 (1H, m); 3.35 , (1H,m); 2.85 , (3H,d); 2.72 , (1H,m).

- 57 -

MS TOF 415 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 11.65 min.

Example 104.

5 **3-Amidino-benzoyl-D-phenylglycine-N-methyl-1-naphthylmethylethylamide**

¹H nmr (DMSO) 8.35 (1H, s); 8.30 (1H, d); 8.10 (1H, d); 7.85 (5H, m); 7.60 (4H, m); 7.45 (4H, m); 6.20 (1H, s); 5.15 (2H, m); 2.91 (3H, s). MS TOF 451 (M+1⁺). Hplc
10 (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 12.53 min.

Example 105.

15 **3-Amidino-benzoyl-D-phenylglycine-S-1-naphthylethylamide**
¹H nmr (DMSO) 8.25 (1H, s); 8.19 (1H, d); 8.15 (2H, m); 7.95 (2H, m); 7.85 (1H, d); 7.70 (1H, t); 7.55 (5H, m); 7.45 (3H, m); 5.75 (1H, s); 5.70 (1H, m); 1.45 (3H, d). MS TOF 451 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 12.21 min.

20

Example 106.

3-Amidino-benzoyl-D-phenylglycine-1-S-cyclohexylethylamide

25 ¹H nmr (DMSO) 8.32 (1H, s); 8.20 (1H, d); 7.95 (1H, d); 7.70 (1H, t); 7.50 (2H, m); 7.30 (3H, m); 5.72 (1H, s); 3.55 (1H, m); 0.95-1.80 (11H, m); 0.9 (3H, d). MS TOF 407 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 12.11 min.

30

Example 107.

3-Amidino-benzoyl-D-phenylglycine-+/- (2-methyl)cyclohexylamide

35 ¹H nmr (DMSO) mixture of isomers (major isomer) 8.29 (1H, s); 8.18 (1H, d); 7.90 (1H, d); 7.70 (1H, t); 7.50 (2H, m); 7.30 (3H, m); 5.70 (1H, s); 0.9-1.8 (8H, m); 0.55 (3H, d). MS TOF 3.93 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 11.45 min.

- 58 -

Example 108.**3-Amidino-benzoyl-D-phenylglycine-exo-2-norbornanamide**

¹H nmr (CD₃CN) 8.25 (1H, s); 8.17 (1H, d); 7.90 (1H, d);
7.68 (1H, t); 7.50 (2H, m); 7.30 (3H, m); 5.55 (1H, s);
5 3.62 (1H, m); 1.05-1.80 (10H, m). MS TOF 391 (M+1⁺).
Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA)
rt 11.18 min.

Example 109.

10 **3-Amidino-benzoyl-D-phenylglycine-4-
chlorobenzhydramide**

¹H nmr (DMSO) 8.35 (1H, s); 8.29 (1H, d); 8.03 (1H, d);
7.79 (1H, t); 7.62 (2H, d); 7.41 (10H, s); 7.18 (2H, d);
6.23 (1H, s); 6.00 (1H, s). MS TOF 498 (M+1⁺). Hplc
15 (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt
13.50 min.

Example 110.

20 **3-Amidino-benzoyl-D-phenylglycine-1,1-
diphenylmethanamide**

¹H nmr (DMSO) 8.23 (1H, s); 8.17 (1H, d); 7.95 (1H, d);
7.70 (1H, t); 7.52 (2H, m); 7.35 (11H, s); 7.05 (2H, m);
6.10 (1H, s); 5.75 (1H, s). MS TOF 464 (M+1⁺). Hplc
(Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt
25 12.04 min.

Example 111.**3-Amidino-benzoyl-D-phenylglycine-4-ethylbenzylamide**

¹H nmr (DMSO) 8.35 (1H, s); 8.25 (1H, d); 7.95 (1H, d);
30 7.70 (1H, t); 7.62 (2H, d); 7.41 (5H, m); 7.21 (1H, t);
6.90 (1H, d); 5.90 (1H, s); 2.60 (2H, q); 1.20 (3H, t).
MS TOF 401 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 12.03 min.

35

- 59 -

Example 112.**3-Amidino-benzoyl-D-phenylglycine-2-tertiary butyl anilide**

¹H nmr (DMSO) 8.21 (1H, s); 8.12 (1H, d); 7.85 (1H, d);
5 7.60 (1H, t); 7.52 (2H, d); 7.25 (4H, m); 7.10 , (2H, m);
6.90 , (1H, m); 5.90 , (1H, s); 1.00 , (9H, s). MS TOF 429
(M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 12.05 min.

10 Example 113.**3-Amidino-benzoyl-D-phenylglycine-3-(1-hydroxyethyl) anilide**

¹H nmr (DMSO) 8.45 (1H, s); 8.35 (1H, d); 8.08 (1H, d);
7.88 (1H, t); 7.75 (3H, m); 7.55 (4H, m); 7.38 , (1H, t);
15 7.20 , (1H, d); 6.02 , (1H, s); 4.80 , (1H, m); 1.45 , (3H, d).
MS TOF 417 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 9.12 min.

Example 114.**20 3-Amidino-benzoyl-D-phenylglycine-4-(4'-acetylphenyl)piperazinamide**

¹H nmr (CD₃CN) 8.12 (1H, s); 8.05 (1H, d); 7.87 (1H, d);
7.80 , (2H, d); 7.65 (1H, t); 7.41 (5H, m); 6.80 , (2H, d);
6.05 (1H, s); 3.80 (1H, m); 3.65 (2H, m); 3.45 (2H, m);
25 3.35 (1H, m); 3.20 (1H, m); 2.75 (1H, m); 2.45 (3H, s). MS
TOF 485 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1,
water/acetonitrile/TFA) rt 12.96 min.

Example 115.**30 3-Amidino-benzoyl-D-phenylglycine-4-(4'-methoxyphenyl)piperazinamide**

¹H nmr (CD₃CN) 8.15 (1H, s); 8.08 (1H, d); 7.91 (1H, d);
7.68 , (2H, d); 7.45 (1H, t); 7.41 (5H, m); 6.99 , (2H, d);
6.12 (1H, s); 3.95 (3H, m); 3.78 (3H, s); 3.75 (2H, m);
35 3.55 (2H, m); 3.45 (1H, m); 2.92 (1H, m). MS TOF 473
(M+1⁺). Hplc (Jupiter5 C18, Gradient 1,
water/acetonitrile/TFA) rt 10.34 min.

- 60 -

Example 116.**3-Amidino-benzoyl-D-phenylglycine-4-(4'-chlorophenyl)piperazinamide**

¹H nmr (CD₃CN) 8.05 (1H, s); 8.00 (1H, d); 7.87 (1H, d);
5 7.55 (1H, t); 7.31 (5H, m); 7.08 (2H, d); 6.75 (2H, d);
5.95 (1H, s); 3.70 (1H, m); 3.55 (2H, m); 3.45 (1H, m);
3.12 (1H, m); 3.00 (1H, m); 2.85 (1H, m); 2.35 (1H, m). MS
TOF 477 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1,
water/acetonitrile/TFA) rt 17.80 min.

10

Example 117.**3-Amidino-benzoyl-D-phenylglycine-4-phenylpiperazinamide**

¹H nmr (DMSO) 8.28 (1H, s); 8.21 (1H, d); 7.94 (1H, d);
7.71 (1H, t); 7.52 (2H, d); 7.35 (3H, m); 7.21 (2H, m);
15 6.89 (2H, d); 6.80 (1H, t); 6.15 (1H, s); 3.75 (3H, m);
3.55 (1H, m); 3.15 (2H, m); 3.00 (1H, m); 2.78 (1H, m). MS
TOF 442 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 9.99 min.

20

Example 118.**3-Amidino-benzoyl-D-phenylglycine-4-toluenesulphonyl piperazinamide**

¹H nmr (DMSO) 8.21 (1H, s); 8.11 (1H, d); 7.91 (1H, d);
7.69 (1H, t); 7.50 (4H, m); 7.35 (2H, m); 7.31 (3H, m);
25 6.05 (1H, s); 3.60 (3H, m); 3.45 (2H, m); 2.90 (1H, m);
2.78 (1H, m); 2.70 (1H, m); 2.45 (3H, s); 2.25 (1H, m). MS
TOF 521 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 10.76 min.

30

Example 119.**3-Amidino-benzoyl-D-phenylglycine-4-benzoyl piperazinamide**

¹H nmr (CD₃CN) 7.95 (1H, s); 7.88 (1H, d); 7.70 (1H, d);
7.45 (1H, t); 7.20 (10H, m); 5.85 (1H, s, broad); 2.50-
35 3.60 (8H, m, broad signals). MS TOF 470 (M+1⁺). Hplc
(Jupiter5 C18, Gradient 1, water/acetonitrile/TFA) rt
10.78 min.

- 61 -

Example 120.**3-Amidino-benzoyl-D-phenylglycine piperidinamide**

¹H nmr (DMSO) 8.20 (1H, s); 8.10 (1H, d); 7.86 (1H, d);
7.65 (1H, t); 7.45 (5H, m); 6.02 (1H, s); 3.70 (1H, m);
5 3.40 (3H, m); 1.53 (3H, m); 1.41 (2H, m); 0.93 (1H, m).
MS TOF 365 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 9.29 min.

Example 121.

10 **3-Amidino-benzoyl-D-phenylglycine-(1S, 2S, 3S, 5R)-
isopinocampyl amide**

¹H nmr (DMSO) 8.30 (1H, s); 8.21 (1H, d); 7.98 (1H, d);
7.71 (1H, t); 7.51 (2H, m); 7.35 (3H, m); 5.74 (1H, s);
4.05 (1H, m); 2.35 (2H, m); 1.90 (1H, m); 1.72 (2H, m);
15 1.60 (1H, m); 1.21 (3H, s); 1.01 (1H, d); 0.97 (3H, s); 0.83
(3H, d). MS TOF 433 (M+1⁺). Hplc (Jupiter5 C18, Gradient
3, water/acetonitrile/TFA) rt 13.32 min.

Example 122.

20 **3-Amidino-benzoyl-D-phenylglycine-3-phenylanilide**

¹H nmr (DMSO) 8.31 (1H, s); 8.24 (1H, d); 7.95 (2H, m);
7.53 (1H, t); 7.60 (5H, m); 7.4 (8H, m); 5.85 (1H, s).
MS TOF 449 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 12.42 min.

25

Example 123.

**3-Amidino-benzoyl-D-phenylglycine-3-(O-benzoyl
hydroxymethyl)anilide**

¹H nmr (DMSO) 8.40 (1H, d); 8.33 (1H, s); 8.31 (1H, d);
30 7.97 (2H, d); 7.75 (1H, t); 7.64 (1H, m); 7.56 (2H, d); 7.40
(8H, m); 7.21 (1H, t); 5.75 (1H, s); 5.20 (2H, s). MS TOF
507 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 13.53 min.

35

- 62 -

Example 124.**3-Amidino-benzoyl-D-phenylglycine-4-(4-chlorobenzoyl) piperidinamide**

¹H nmr (DMSO) Mixture of rotomers 8.08 (2H, m); 7.80 (3H, m); 7.55 (1H, t); 7.45 (2H, m); 7.25 (5H, m); 5.92 (1H, s); 4.25 (1H, m); 3.80 (1H, m); 3.55 (1H, m); 2.90 (1H, m); 2.68 (1H, m); 1.60 (1H, m); 1.55 (1H, m); 1.30 (1H, m); 0.83 (1H, m). MS TOF 504 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 12.21 min.

Example 125.**3-Amidino-benzoyl-D-phenylglycine-4-(4-fluorobenzoyl) piperidinamide**

¹H nmr (DMSO) Mixture of rotomers 8.40 (2H, m); 8.22 (2H, m); 8.10 (1H, d); 7.88 (1H, t); 7.64 (2H, m); 7.54 (5H, m); 6.30 (1H, s); 4.65 (1H, m); 4.20 (1H, m); 3.82 (1H, m); 3.10 (2H, m); 1.98 (1H, m); 1.70 (1H, m); 1.40 (1H, m); 0.83 (1H, m). MS TOF 487 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 11.37 min.

Example 126.**3-Amidino-benzoyl-D-phenylglycine-4-benzyl piperidinamide**

¹H nmr (DMSO) Mixture of rotomers 7.98 (2H, m); 7.68 (1H, m); 7.45 (1H, m); 6.60-7.25 (10H, m); 5.86 (1H, s); 4.15 (1H, m); 3.68 (1H, m); 2.63 (1H, m); 2.45 (1H, m); 2.21 (3H, m); 1.50 (1H, m); 1.33 (1H, m); 1.03 (1H, m); 0.45 (1H, m). MS TOF 455 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 13.18 min.

Example 127.**3-Amidino-benzoyl-D-phenylglycine-4-D-2-naphthylalanine pyrrolidinamide**

¹H nmr (DMSO) 8.19 (1H, s); 8.11 (1H, d); 7.92 (1H, d); 7.75 (5H, m); 7.41 (5H, m); 7.30 (3H, m); 5.81 (1H, s); 4.80 (1H, m); 3.20 (4H, m); 3.00 (2H, m); 1.55 (4H, m). MS TOF 548 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,

- 63 -

water/acetonitrile/TFA) rt 11.13 min.

Example 128.

3-Amidino-benzoyl-DL-1-naphthylglyciny-D-2-naphthylalanine pyrrolidinamide

¹H nmr (DMSO) mixture of isomers 8.25 (1H, s); 8.20 (1H, d); 8.15 (2H, m); 8.05 (1H, d); 6.90-8.00 complex multiplet (13H, m); 6.50 (1H, m); 4.90 (1H, m); 2.90-3.50 broad, partially covered by solvent (H, m); 1.70 (4H, m). MS TOF 598 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 12.18 and 12.37 min.

Example 129.

3-Amidino-benzoyl-DL-1-naphthylglycine-4-methylbenzylamine

¹H nmr (DMSO) mixture of isomers 8.25 (1H, s); 8.20 (1H, d); 8.10 (1H, m); 7.90 (3H, m); 7.70 (1H, t); 7.55 (4H, m); 7.06 (4H, m); 6.45 (1H, s); 4.30 (2H, m); 2.25 (3H, s). MS TOF 451 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 12.30 min.

Example 130.

3-Amidino-benzoyl-DL-1-naphthylglycine-4-benzoyl piperidinamide

¹H nmr (DMSO) mixture of isomers 8.15 (2H, m); 8.00 (1H, d); 7.85 (5H, m); 7.45 (8H, m); 6.70 (1H, s); 4.50 (1H, m); 3.15 (1H, m); 2.80 (3H, m); 1.60 (2H, m); 1.42 (1H, m); 0.45 (1H, m). MS TOF 519 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 12.67 min.

Example 131.

3-Amidino-benzoyl-DL-1-naphthylglycine-3-(1-hydroxyethyl)anilide

¹H nmr () mixture of isomers 8.10 (1H, s); 8.00 (1H, d); 7.80 (3H, m); 7.50 (7H, m); 7.30 (1H, m); 7.20 (1H, m); 7.00 (1H, m); 6.45 (1H, s); 4.65 (1H, m); 1.20 (3H, s). MS TOF 467 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,

- 64 -

water/acetonitrile/TFA) rt 10.67 min.

Example 132.

3-Amidino-benzoyl-D-phenylglycine morpholinamide

5 ¹H nmr (DMSO) 8.25 (1H, s); 8.18 (1H, d); 7.91 (1H, d);
7.68 (1H, t); 7.40 (5H, m); 6.09 (1H, s); 5.45 (4H, m);
3.63 (4H, m). MS TOF 367 (M+1⁺). Hplc (Jupiter5 C18,
Gradient 3, water/acetonitrile/TFA) rt 5.53 min.

10 **Example 133.**

3-Amidino-benzoyl-R,S-3-amino-3-phenylpropionic-1-adamantylamide

¹H nmr (CD₃CN) 8.03 (1H, s); 7.97 (1H, d); 7.75 (1H, d);
7.55 (1H, t); 7.20 (5H, m); 5.25 (1H, t); 2.50-1.70 (15H,
15 m). MS TOF 446 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 11.56 min.

Example 134.

3-Amidino-benzoyl-DL-1-naphthylglycine-(Z,E)-(2-methyl)cyclohexylamide

20 Mixture of isomers. MS TOF 443 (M+1⁺). Hplc (Jupiter5
C18, Gradient 3, water/acetonitrile/TFA) rt 12.57 and
12.75 min.

25 **Prepared by Method 5**

Example 135.

3-Amidino-benzoyl-D-phenylglycine cyclopropylmethylethylamide

MS TOF 351 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
30 water/acetonitrile/TFA) rt 8.57 min.

Example 136.

3-Amidino-benzoyl-D-phenylglycine-2-indanylamide

MS TOF 413 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
35 water/acetonitrile/TFA) rt 11.13 min.

- 65 -

Example 137.

3-Amidino-benzoyl-D-phenylglycine-3-methylbutylamide
MS TOF 367 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 10.81 min.

5

Example 138.

3-Amidino-benzoyl-D-phenylglycine-4-fluorobenzylamide
MS TOF 405 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 10.41 min.

10

Example 139.

3-Amidino-benzoyl-D-phenylglycine trans-2-phenylcyclopropylamide
MS TOF 413 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 11.25 min.

15

Prepared by Method 6**Example 140.**

20

**3-Amidino-benzoyl-D-phenylglycine-2-(4-N-acetyl
piperidinyl)ethyl ester**

¹H nmr (DMSO) 8.13 (1H, s); 8.07 (1H, d); 7.92 (1H, d);
7.69 (2H, d); 7.45 (5H, m); 5.63 (1H, s); 4.25 (2H, m);
3.65 (1H, m); 3.45 (3H, m); 2.80 (1H, m); 3.35 (1H, m); 2.00
(3H, s); 1.50 (3H, m); 1.30 (1H, m); 0.95 (1H, m). MS TOF
451 (M+1⁺). Hplc (Jupiter5 C18, Gradient 2,
water/acetonitrile/TFA) rt 19.00 min.

25

Method 7

30

By solution phase strategy: Typically a Boc-amino
alcohol with TEA is treated with an acid chloride to
give an ester. Removal of the Boc protecting group with
TFA followed by further extension of the compounds as in
Method 4.

35

- 66 -

Example 141.

{R}-{[N-(3-amidino)benzoyl]amino}-2-phenylethyl-4-(methoxy)benzoate.

5 To Boc-(D)-phenylglycinol (237mg, 1mmol, 1eq) and TEA(153 μ l, 1.1mmol, 1eq) dry DCM (5ml) was added p-anisoyl chloride (188mg, 1.1mmol, 1.1eq). The reaction was stirred overnight. Ethylacetate (30ml) was added and the organic layer was washed successively with 2 x
10 10ml of 10%HCl, NaHCO₃ solution, finally brine. Dried over magnesium sulphate, filtered and evaporated. to give the crude product (410mg). After treatment with TFA to remove the Boc group the compound were extended as in Method 4.

15

¹H nmr (DMSO) 8.25 (1H, s); 8.21 (1H, d); 7.95 (1H, d); 7.85 (2H,d); 7.75 (1H, t); 7.45 (5H, m); 7.05 (2H,d); 5.55 (1H, q); 4.55 (2H, d); 3.85 (3H, s). MS TOF 418 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 11.01 min.

20

By analogous methods the following were prepared

Example 142.

25 **R-2-{[N-(3-amidino)benzoyl]amino}-2-phenylethyl-4-(methyl)benzoate**

¹H nmr (DMSO) 8.25 (1H, s); 8.21 (1H, d); 7.95 (1H, d); 7.80 (2H,d); 7.75 (1H, t); 7.50 (2H,d); 7.34 (5H, m); 5.55 (1H, q); 4.55 (2H, d); 3.35 (3H, s). MS TOF 402 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 11.87 min.

30

Example 143.

35 **R-2-{[N-(3-amidino)benzoyl]amino}-2-phenylethyl-4-(trifluoromethoxy)benzoate**

¹H nmr (DMSO) 8.45 (1H, s); 8.40 (1H, d); 8.25 (2H,d); 8.15 (1H, d); 7.95 (1H, t); 7.65 (7H, m); 5.78 (1H, q);

- 67 -

4.84 (2H, d). MS TOF 472 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 12.85 min.

Method 8

5

By solution phase strategy: Typically a Boc-amino alcohol and a carboxylic acid are treated with (4-dimethylamino) phenyldiphenyl phosphine and diethylazodicarboxylate to give an ester. Removal of the Boc protecting group with TFA followed by further extension of the compounds as in Method 4.

10

Example 144.

{R}-[N-(3-amidino)benzoyl]amino}-2-phenylethyl-4-(acetimido)benzoate

15

A dry solution of diethylazodicarboxylate (158 μ l, 1mmol, 1eq) in THF (5ml), was added dropwise to an anhydrous solution of (4-dimethylamino)phenyldiphenyl phosphine (306mg, 1eq, 1mmol), Boc-(D)-phenylglycinol (356mg, 1.5eq, 1mmol) and 4-acetimidobenzoic acid (179mg, 1mmol, 1eq) in THF (5ml) at -78°C. The reaction allowed to warm to room temperature and stirred overnight. Reaction monitored by t.l.c, SiO₂ / 75% ethylacetate in hexane indicating formation of product. R_f = 0.5. The solution was evaporated to remove THF and taken up into ethyl acetate(30ml) washed with 10% HCl solution, then brine. The solution was dried, filtered and evaporated to dryness. Crude yield ~650mg. Purification by column chromatography gave a yield of product of 200mg, (50%). After treatment with TFA to remove the Boc group the compound were extended as in Method 4.

20

25

30

¹H nmr (DMSO) 8.48 (1H, s); 8.54 (1H, d); 8.15 (1H, d); 8.05 (2H, d); 7.95 (3H, m); 7.75 (2H, m); 7.55 (3H, m); 5.70 (1H, m); 4.70 (2H, m); 2.27 (3H, s). MS TOF 445 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,

35

- 68 -

water/acetonitrile/TFA) rt 9.38 min.

Example 145.

R-2-{ [N-(3-amidino)benzoyl]amino}-2-phenylethyl-4-(methylsulphonyl)benzoate

¹H nmr (DMSO) 8.25 (1H, s); 8.21 (1H, d); 8.10 (4H, q); 7.95 (1H, d); 7.75 (1H, t); 7.45 (5H, m); 5.60 (1H, q); 4.65 (2H, d); 2.08 (3H, s). MS TOF 466 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 9.75 min.

Example 146.

R-2-{ [N-(3-amidino)benzoyl]amino}-2-phenylethyl-4-acetylbenzoate

¹H nmr (DMSO) 8.03 (1H, s); 8.00 (1H, d); 7.83 (4H, m); 7.73 (1H, d); 7.53 (1H, t); 7.25 (5H, m); 5.38 (1H, q); 4.41 (2H, d); 2.42 (3H, s). MS TOF 430 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 10.46 min.

Method 9

By solution phase strategy: Typically an activated Boc-amino acid was converted to the amide with ammonia and then to the thioamide using Lawessons reagent. Reaction of the thioamide with an acyl bromide to give a Boc protected thiazole. Deprotection of the thiazole with TFA allows further extension of the compound as in Method 4.

Example 147.

2-[(N-3-amidinobenzamido)-1-phenyl]methyl-4-phenylthiazole

To a solution of Boc-D-phenylglycine (875mg 3.5mmole) in a 1:1 mixture of DMF and DCM (40ml) was added 1-hydroxybenzotriazole (520mg 1.1equiv.) and DIPCI (602ml

- 69 -

1.1equiv.) and the mixture stirred for 30min at rt. Ammonia gas was bubbled in and the mixture left overnight before diluting with ethyl acetate and washing with 10% hydrochloric acid and sat. sodium bicarbonate.

5 The organic solution was dried (magnesium sulphate) and evaporated. Flash chromatography (silica gel DCM/ethylacetate 0-50%) gave the amide 770mg 86%. To a solution of Boc-D-phenylglycine amide (740mg 2.96mmole) in THF 25ml was added Lawesson's reagent

10 (1.2g) and the mixture stirred overnight. The solvent was evaporated off under reduced pressure and the residue purified by flash chromatography (silica gel - hexane/ethyl acetate 10 to 30%) to give the thioamide 671mg.

15 The thioamide (650mg) was dissolved in acetone (20ml) and phenacyl bromide (486mg 1 equiv) stirred for 30min and then diluted with chloroform/aqueous sod. bicarbonate (20ml each), separated, dried (magnesium sulphate) and evaporated. The residue was dissolved in

20 DCM (20ml) and treated with pyridine (350ml) and trifluoroacetic anhydride (360ml). After 150min the solvent was removed under reduced pressure and the residues redissolved in DCM and washed with sat. sod. bicarbonate. Purification by flash chromatography

25 (silica gel - hexane/ethyl acetate 10 to 30%) gave the thiazole intermediate 687mg. Deprotection of the amine was carried out using 50% TFA in DCM (30min), evaporation of solvent under reduced pressure, dissolving in DCM, washing with sat sod. bicarbonate, drying (magnesium sulphate) and

30 evaporation. Coupling of the thiazole intermediate to 3-amidino benzoic acid was carried out using the standard method Method 4 to give 2-[(N-3-amidinobenzamido)-1-phenyl]methyl-4-phenylthiazole TFA salt.

35

¹HNMR (d₆DMSO) 9.9 (1H, d); 9.48 (1H, s); 9.09 (1H, s);

- 70 -

8.40 (1H, s); 8.37 (1H, d); 8.16 (1H, s); 7.98 (3H, m);
7.80 (2H, t); 7.63 (2H, d); 7.5 (6H, m); 6.80 (1H, d). MS
TOF 413 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 11.8 min.

5

Method 10

By solution phase: Typically a compound prepared as
Method 4 is further reacted with ethyl chloroformate to
10 give the 3-Ethoxycarbonyl amidino compound

Example 148.

**3-(Ethoxycarbonyl)-amidinobenzoyl-D-phenylglycine-4-
methylbenzyl amide.**

15

To a solution of 3-Amidinobenzoyl-(D)-phenylglycine-4-
methylbenzyl amide (125mg, 0.31mmole) in DCM (20ml) was
added DIPEA (163ml 0.94mmole) and then over 2min. ethyl
chloroformate 33ml 0.34mmole). The reaction was stirred
20 overnight and the solvent removed under reduced
pressure. The residue was dissolved in ethylacetate and
washed with water (x3), dried (magnesium sulphate) and
evaporated to dryness. Flash chromatography (silica gel,
DCM/Ethylacetate 0-100%) gave the title compound 98mg

25

¹HNMR (d₆DMSO) 9.1 (2H, bs); 8.93 (1H, d); 8.82 (1H, t);
8.46 (1H, s); 8.08 (2H, t); 7.53 (3H, m); 7.36 (3H, m);
7.08 (4H, s); 5.76 (1H, d); 4.23 (2H, d); 4.07 (2H, q);
2.25 (3H, s); 1.22 (3H, t). MS TOF 473 (M+1⁺). Hplc
30 (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt
11.06 min.

By an analogous method

35

- 71 -

Example 149**3-(Ethoxycarbonylamidino)-D-phenylglycine-4-(4-chlorophenyl)-piperazinamide**

¹HNMR (d₆DMSO) 9.01 (3H, bs+s); 8.40 (1H, s); 8.08 (2H, m); 7.52 (3H, m); 7.40 (3H, m); 7.25 (2H, d); 6.88 (2H, d); 6.15 (1H, d); 4.05 (2H, d); 3.70 (4H, m); 3.01 (4H, m); 1.20 (3H, t). MS TOF 549 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 12.48 min.

Method 11

By solution phase strategy: Typically 3-cyanobenzoic acid is activated with carbonyl-diimidazole and reacted with R-phenylglycinol to give 3-cyanobenzoylphenylglycinol. Reaction with a trichloroacetimidate and BF₃ etherate, (4), gives an ether. Reaction of the cyano group with HCl in ethanol followed by ammonia in ethanol gives the amidino compound.

Example 150.**3-Amidinobenzoyl-2-(4-methylbenzyloxy)-1-phenylethylamide**

3-Cyanobenzoic acid was condensed with (R)-phenylglycinol using N,N'-carbonyl-diimidazole as activator to give 3-cyanobenzoylphenylglycinol. ¹HNMR (CDCl₃) 8.05 (2H, d); 7.55 (1H, t); 7.4 (8H, m); 5.17 (2H, bs); 4.50 (2H, bs).

3-Cyanobenzoylphenylglycinol in 1:1 DCM: cyclohexanol (50ml) was treated with 4-methylbenzyl trichloroacetimidate and BF₃ etherate and the crude product purified by flash chromatography to give 3-cyanobenzoyl-2-(4-methylbenzyloxy)-1-phenylethylamide which was then taken up in ethanol and saturated with HCl gas and left overnight at room temperature. The solvent and excess HCl gas were removed under reduced

- 72 -

pressure and the residue redissolved in ethanol and the solution saturated with ammonia gas and left overnight. The solvent and excess ammonia were removed under reduced pressure and the residue treated with aqueous acetonitrile and trifluoroacetic acid. Preparative hplc gave the title compound approx 80% pure by hplc.

¹HNMR (d₆DMSO) 9.1 (5H, bs+d); 8.22 (2H, s+d); 7.90 (1H, d); 7.74 (1H, t); 7.35 (5H, m); 7.10 (4H, Abq); 5.35 (1H, q); 4.50 (2H, s); 3.70 (2H, m); 2.25 (3H, s). MS TOF 389 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 11.79 min.

Miscellaneous Examples

Example 151.

N-(3-Amidinobenzoyl)-2-amino-2(R)-phenylethoxycarbonyl-4-methylaniline

To a solution of Boc-(R)-phenylglycinol (500mg 3.65mmole) in DMF (20ml) was added 4-methylphenylisocyanate (500ml 4.0mmole) and the mixture warmed to 50°C before standing at room temperature overnight. Dilution with ethyl acetate and washing with sat. sodium bicarbonate gave after drying (magnesium sulphate) and evaporation of solvent the crude intermediate which was purified by flash chromatography (silica gel, EtOAc 0-20% in DCM)

Deprotection of the amine was carried out using 50% TFA in DCM (30min), evaporation of solvent under reduced pressure, dissolving in DCM, washing with sat sod. bicarbonate, drying (magnesium sulphate) and evaporation.

Coupling of the thiazole intermediate to m-benzamidine was carried out using the standard method 4 to give N-(3-Amidinobenzoyl)-2-amino-2(R)-phenylethoxycarbonyl-4-methylaniline.

- 73 -

¹H NMR (d₆DMSO) 8.2 (2H, m); 7.90 (1H, d); 7.72 (1H, t); 7.44 (2H, d); 7.35 (5H, m); 7.08 (2H, d); 5.35 (1H, m); 4.33 (2H, m); 2.25 (3H, s). MS TOF 417 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 11.54 min.

Example 152.**3-Amidinobenzyl-D-phenylglycine-cis/trans-4-aminomethyl cyclohexylmethyleamide**

10

Bis-1,4-aminomethylcyclohexane 2-chlorotrityl resin (0.2 mmol) (as prepared by method 1) was treated with Fmoc-D-Phenylglycine (0.5 mmol, 187mg), DMF (2.5ml), TBTU in DMF (1.25ml of a 450mM solution) and DIPEA in DMF (1.25ml of a 900 mM solution). The mixture was agitated with nitrogen for 2 hours. Deprotection and washing as in method 1. The resin was then treated with 3-cyanobenzaldehyde (0.5mmol, 66mg) in DMF agitated overnight. The resin was washed with DMF then with Dry THF to remove all DMF. Sodium cyanoborohydride (1mmol, 63mg) in THF and acetic acid (100μl) was then added to the resin and agitated for 4 hours. The resin was then washed with THF, the DMF and finally DCM. The resin was then treated with TFA containing 5% TES for 30 mins. The resin was then filtered off and the TFA evaporated then held under high vacuum to dry the product. The product was then taken up into saturated hydrogen chloride gas in ethanol (20ml.) overnight. The ethanol was then evaporated and the product treated with saturated ammonia gas in ethanol (20ml.) overnight. The ethanol was then evaporated and the product purified by preparative hplc.

¹H nmr (CD₃CN) Mixture of cyclohexyl cis and trans isomers 7.85 (2H, m); 7.79 (1H, d); 7.66 (1H, t); 7.50 (5H, m); 4.97 (1H, s); 4.22 (2H, s); 3.00 (2H, m); 2.78 (2H, m); 1.48 (7H, m); 0.86 (3H, m) MS TOF 407 (M+1⁺).

- 74 -

Hplc (Jupiter5 C18, Gradient 2, water/acetonitrile/TFA)
rt 9.78 min.

Example 153.**5 4-Methylphenylacetyl-L-phenylglycine-3-amidino anilide**

Boc-L-phenylglycine (0.6mmol, 150mg), HATU (0.6mmol, 230mg) and DIPEA (1.2mmol., 220 μ l) were dissolved in DMF (5ml) and stirred for 15mins. 3-Aminobenzonitrile (0.6mmol., 70mg) was added and the mixture stirred overnight. The DMF was then evaporated and the mixture taken into ethylacetate (50ml) and washed with water, sodium carbonate solution, water 10% hydrochloric acid and water. The ethylacetate was evaporated to give the product Boc-L-phenylglycine-3-cyano anilide. The product was treated with TFA (50ml) for 30mins. The TFA was evaporated to dryness. m-Tolylacetic acid (0.7mmol., 100mg), HATU (0.7mmol., 250mg) and DIPEA (1.4mmol., 250 μ l) were dissolved in DMF (5ml) and stirred for 15mins. This was then added to the above product with DIPEA (0.6mmol., 110 μ l) and stirred overnight. The DMF was evaporated and the mixture taken into ethylacetate (50ml) and washed as above. The ethyl acetate was then evaporated and the product treated with hydrogen chloride gas in ethanol and ammonia gas in ethanol as above. The product was purified by preparative hplc.

¹HNMR (d₄ MeOH) 8.23 (1H, s); 7.90 (1H, d); 7.72 (7H, m); 7.44 (1H, t); 7.29 (3H, m); 5.75 (1H, m); 3.80 (2H, s); 2.50 (3H, s). MS TOF 401 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA) rt 14.59 min.

Compounds synthesised by Method 4

35

- 75 -

Example 154.**3-Amidino-benzoyl-D-phenylglycine-4-(4'-nitrophenyl)
piperazinamide**

5 ¹H nmr (CD₃CN) 8.20 (1H, s); 8.05 (3H, m); 7.85 (1H, d);
7.70 (1H, t); 7.40 (5H, m); 6.82 (2H, d); 6.05 (1H, s);
3.75 (1H, m); 3.65 (2H, m); 3.45 (3H, m); 3.30 (1H, m); 2.85
(1H, m). MS TOF 486 (M+1⁺). Hplc (Jupiter5 C18, Gradient
1, water/acetonitrile/TFA) rt 11.65 min.

10

Example 155.**3-Amidino-benzoyl-D-phenylglycine-4-(4'-aminophenyl)
piperazinamide**

15 ¹H nmr (CD₃CN) 8.00 (1H, s); 7.85 (1H, d); 7.65 (1H, d);
7.45 (1H, t); 7.20 (5H, m); 6.95 (2H, d); 6.70 (2H, d);
5.85 (1H, s); 3.55 (1H, m); 3.30 (1H, m); 2.95 (3H, m);
2.80 (1H, m); 2.40 (1H, m). MS TOF 458 (M+1⁺). Hplc
(Jupiter5 C18, Gradient 2, water/acetonitrile/TFA) rt
20 12.57 min.

Example 156.**3-Amidino-benzoyl-D-phenylglycine-4-(4'-fluorophenyl)
piperazinamide**

25

¹H nmr (CD₃OD) 8.25 (1H, s); 8.20 (1H, d); 7.95 (1H, d);
7.75 (1H, t); 7.45 (5H, m); 7.10 (4H, m); 6.20 (1H, s);
3.90 (3H, m); 3.70 (1H, m); 3.35 (2H, m); 3.15 (1H, m); 2.65
(1H, m). MS TOF 460 (M+1⁺). Hplc (Jupiter5 C18, Gradient
30 1, water/acetonitrile/TFA) rt 14.88 min.

Example 157.**3-Amidino-benzoyl-D-phenylglycine-4-(4'-pyridyl)
piperazinamide**

35

¹H nmr (CD₃CN) 8.05 (1H, s); 7.95 (1H, d); 7.90 (2H, d);
7.80 (1H, d); 7.55 (1H, t); 7.30 (5H, m); 6.80 (2H, d);

- 76 -

6.00 (1H, s); 3.70 (3H,m); 3.50 (3H,m); 3.40 (1H,m);
3.15 (1H,m). MS TOF 444 (M+1⁺). Hplc (Jupiter5 C18,
Gradient 2, water/acetonitrile/TFA) rt 11.54 min.

5 **Example 158.**

**3-Amidino-benzoyl-D-phenylglycine-4-(2'-pyridyl)
piperazinamide**

¹H nmr (CD₃OD) 8.30 (1H, s); 8.20 (1H, d); 8.10 (1H, m);
10 7.95 , (2H,m); 7.75 (1H, t); 7.50 (5H, m); 7.30 , (2H,d);
7.15 (1H, t); 6.20 (1H, s); 3.95 (3H,m); 3.75 (4H,m);
3.40 (1H,m). MS TOF 444 (M+1⁺). Hplc (Jupiter5 C18,
Gradient 2, water/acetonitrile/TFA) rt 11.67 min.

15 **Example 159.**

3-Amidino-benzoyl-D-phenylglycine-4-benzylpiperazinamide

¹H nmr (CD₃CN) 8.15 (1H, s); 8.05 (1H, d); 7.90 (1H, d);
7.65 (1H, t); 7.40 (10H, m); 6.05 (1H, s); 4.20 (2H,s);
20 3.20 (8H,v. broad m). MS TOF 457 (M+1⁺). Hplc (Jupiter5
C18, Gradient 1, water/acetonitrile/TFA) rt 9.52 min.

Example 160.

25 **3-Amidino-benzoyl-D-phenylglycine-4-(3-chlorophenyl)
piperazinamide**

¹H nmr (CD₃CN) 8.17 (1H, s); 8.07 (1H, d); 7.90 (1H, d);
7.65 (1H, t); 7.42 (5H, m); 7.15 (1H, m); 6.80 (3H,m);
6.08 (1H, s); 3.85 (1H,m); 3.55 (3H,m); 3.25 (1H,m);
30 3.10 (1H,m); 2.95 (1H,m); 2.40 (1H,m). MS TOF 477
(M+1⁺). Hplc (Jupiter5 C18, Gradient 1,
water/acetonitrile/TFA) rt 15.39 min.

- 77 -

Example 161.**3-Amidino-benzoyl-D-phenylglycine-4-(2-chlorophenyl)
piperazinamide**

5 ¹H nmr (CD₃CN) 8.15 (1H, s); 8.07 (1H, d); 7.90 (1H, d);
7.65(1H, t); 7.42 (5H, m); 7.20 (1H, m); 6.95 (3H,m);
6.05 (1H, s); 3.80 (1H,m); 3.65 (2H,m); 3.45 (1H,m); 3.
00(1H,m); 2.85 (2H,m); 2.40 (1H,m). MS TOF 477 (M+1⁺).
Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA)
10 rt 15.02 min.

Example 162.**3-Amidino-benzoyl-D-phenylglycine-tetrahydroisoquinolin-
2-amide**

15 ¹H nmr (CD₃CN) 8.05 (1H, s); 8.00 (1H, d); 7.70 (1H, d);
7.45(1H, t); 7.25(2H,m); 7.15 (3H, m); 6.90 , (4H,m);
6.00 (1H, s); 4.45 (2H,m); 3.45 (2H,m); 2.55 (2H,m).
MS TOF 413 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
20 water/acetonitrile/TFA) rt 10.92min.

Example 163.**3-Amidino-benzoyl-D-phenylglycinyln-N-benzyl glycine
ethyl ester**

25 ¹H nmr (CD₃CN) 8.15 (1H, s); 8.10 (1H, d); 7.90 (1H, d);
7.70(1H, t); 7.35 (10H, m); 6.10 (1H, s); 4.65 (2H,m);
4.45 (2H,m); 4.05 (2H,m); 1.15 (3H,m). MS TOF 473
(M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
30 water/acetonitrile/TFA) rt 11.32 min.

Example 164.**3-Amidino-benzoyl-D-phenylglycinyln-1,2,3,4-DL-
tetrahydroisoquinoline-3-dimethylamide**

35 ¹H nmr (CD₃OD) mixture of isomers, major reported. 8.25
(1H, s); 8.20 (1H, d); 7.95 (1H, d); 7.75(1H, t); 7.70 -

- 78 -

7.00 (9H, m); 6.35 (1H, s); 5.20 (1H, m); 4.65 (2H, m);
3.15 (2H, m, obscured by solvent); 3.00 (6H, d). MS TOF
484 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1,
water/acetonitrile/TFA) rt 13.81 min.

5

Compounds synthesised by Method 7

Example 165.

R-2-{[N-(3-amidino)benzoyl]amino}-2-phenylethyl-4-
10 (methyl)benzoate

¹H nmr (DMSO) 8.25 (1H, s); 8.21 (1H, d); 7.95 (1H, d);
7.80 (2H, d); 7.75 (1H, t); 7.50 (2H, d); 7.34 (5H, m);
5.55 (1H, q); 4.55 (2H, d); 3.35 (3H, s). MS TOF 402
15 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 11.87 min.

Example 166.

R-2-{[N-(3-Amidino)benzoyl]amino}-2-phenylethyl-4-
20 dimethylaminobenzoate

¹H nmr (CD₃CN) 8.20 (1H, s); 8.05 (1H, d); 7.90 (1H,
d); 7.75 (2H, d); 7.65 (1H, t); 7.45 (2H, d); 7.35 (3H,
m); 6.70 (2H, d); 5.50 (1H, m); 4.55 (2H, m); 1.95
25 (6H, s). MS TOF 431 (M+1⁺). Hplc (Jupiter5 C18, Gradient
3, water/acetonitrile/TFA) rt 11.11 min.

Example 167.

R-2-{[N-(3-Amidino)benzoyl]amino}-2-phenylethyl-4-
30 aminosulphonylbenzoate

¹H nmr (CD₃CN) 8.15 (1H, s); 8.05 (3H, m); 7.90 (3H, m);
7.65 (1H, t); 7.55 (2H, d); 7.35 (3H, m); 5.50 (1H, m);
4.65 (2H, d). MS TOF 467 (M+1⁺). Hplc (Jupiter5 C18,
35 Gradient 3, water/acetonitrile/TFA) rt 9.80 min.

- 79 -

Example 168.**R-2-{[N-(3-Amidino)benzoyl]amino}-2-phenylethyl benzoate**

¹H nmr (CD₃CN) 8.05 (1H, s); 8.00 (1H, d); 7.80 (2H, m);
5 7.70 (1H, d); 7.55 -7.00 (9H, m); 5.45 (1H, m); 4.55
(2H, m). MS TOF 387 (M+1⁺). Hplc (Jupiter5 C18,
Gradient 3, water/acetonitrile/TFA) rt 10.85 min.

Example 169.

10 **R-2-{[N-(3-Amidino)benzoyl]amino}-2-phenylethyl-4-
acetoxybenzoate**

¹H nmr (CD₃CN) 8.00 (1H, s); 7.95 (1H, d); 7.85 (2H, d);
7.75 (1H, d); 7.55 (1H, t); 7.35 (2H, d); 7.20 (3H, m);
15 7.00 (2H, d); 5.35 (1H, m); 4.55 (2H, m); 2.10 (3H, s).
MS TOF 446 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 10.54 min.

Example 170.

20 **R-2-{[N-(3-Amidino)benzoyl]amino}-2-phenylethyl-4-
isopropylbenzoate**

¹H nmr (CD₃CN) 8.05 (1H, s); 8.00 (1H, d); 7.90 (1H, d);
7.75 (2H, d); 7.55 (1H, t); 7.40 (2H, d); 7.25 (5H, m);
25 5.45 (1H, m); 4.55 (2H, m); 2.85 (1H, m); 1.10 (6H, d). MS
TOF 430 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 12.43 min.

Example 171.

30 **R-2-{[N-(3-Amidino)benzoyl]amino}-2-phenylethyl-4-
hydroxybenzoate**

¹H nmr (CD₃CN) 8.15 (1H, s); 8.05 (3H, m); 7.85 (1H, d);
8.80 (2H, d); 7.65 (1H, t); 7.40 (5H, m); 6.80 (2H, d);
35 5.50 (1H, m); 4.55 (2H, d). MS TOF 404 (M+1⁺). Hplc
(Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt
10.05 min.

- 80 -

The following compounds were synthesised by Method 4 and elaborated as Example 89

Example 172.

5 2-Amino-5-amidino-benzoyl-D-phenylglycine benzylamide

¹H nmr (DMSO) 8.30 (1H, s); 7.80 (1H, d); 7.75 (2H, d); 7.45 (8H, m); 7.05 (1H, d); 5.95 (1H, s); 4.55 (2H, s).

MS TOF 402 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA) rt 10.55 min.

Example 173.

2-Amino-5-amidino-benzoyl-D-phenylglycine-2-methylbenzylamide

¹H nmr (DMSO) 8.15 (1H, s); 7.70 (1H, d); 7.60 (2H, d); 7.30 (7H, m); 6.90 (1H, d); 5.80 (1H, s); 4.30 (2H, s); 2.25 (3H, s). MS TOF 416 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA) rt 10.07 min.

Example 174.

2-Amino-5-amidino-benzoyl-D-phenylglycine-3-methylbenzylamide

¹H nmr (DMSO) 8.15 (1H, s); 7.60 (1H, d); 7.50 (2H, d); 7.35 (4H, m); 7.15 (1H, m); 6.95 (2H, m); 6.80 (1H, d); 5.70 (1H, s); 4.30 (2H, s); 2.25 (3H, s). MS TOF 416 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA) rt 10.94 min.

Example 175.

2-Amino-5-amidino-benzoyl-D-phenylglycine-R- α -methylbenzylamide

¹H nmr (DMSO) 8.15 (1H, s); 7.60 (1H, d); 7.50 (2H, d); 7.35 (5H, m); 7.15 (1H, m); 6.95 (2H, m); 6.80 (1H, d); 5.70 (1H, s); 5.00 (1H, m); 1.45 (3H, d). MS TOF 416

- 81 -

(M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA) rt 10.58 min.

Example 176.

5 **2-Amino-5-amidino-benzoyl-D-phenylglycine-1-naphthylamide**

¹H nmr (DMSO) 8.00 (1H, s); 7.85 (2H,m); 7.75 (1H,d);
7.50 (1H,d); 7.40 (4H,m); 7.25 (5H,m); 6.70 (1H,d);
10 5.60 (1H, s); 4.65 (2H,s). MS TOF 452 (M+1⁺). Hplc
(Jupiter5 C18, Gradient 1, water/acetonitrile/TFA) rt
11.17 min.

Example 177.

15 **2-Amino-5-amidino-benzoyl-D-phenylglycine-2-methylcyclohexylamide**

¹H nmr (DMSO) 8.00 (1H, s); 7.50 (1H, d); 7.40 (2H, m);
7.25 (4H, m); 6.70 (1H,d); 5.55 (1H, s); 3.20 (1H,m);
20 1.7 -0.7 (9H, broad m) 0.70 (3H,d). MS TOF 408 (M+1⁺).
Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA)
rt 10.95 min.

Compounds synthesised by Method 12

25

Example 178.

R-2-{[N-(3-Amidino)benzoyl]amino}-2-phenylethyl-4-methoxybenzamide

30 Typically Boc-(R)-1,2-diamino-1-phenylethane was prepared by the Method of O'Brien, P. et al. *J. Med. Chem.* 37 (1994) 12, 1810-1822. The free amino group was reacted with an acid chloride or an activated acid, all couplings (minimum 120 min.) were carried out in DMF.
35 After an aqueous work up the deprotection of the Boc group was achieved with TFA. Other acid substituents were added as the HOBt or HOAt esters either by

- 82 -

activation with HBTU/HATU, EDC or DIPCI with or without Boc protection of amino groups. The final products were purified by preparative reverse phase Hplc.

5 Boc-(R)-1,2 diamino-1-phenylethane (2g) was dissolved in DCM (80ml) and toluoylchloride (1.2ml.) and TEA (1.1ml.) added. The mixture was stirred at room temperature under argon overnight. The mixture was then washed with 2M
10 sodium hydroxide solution, water, 5% hydrochloric acid and brine, dried over magnesium sulphate, filtered and evaporated. The product was recrystallised from hexane dissolved in DCM (50ml.) and treated with TFA for 2hrs. The solvent was evaporated and the solid triturated with diethylether. Coupling to m-benzamidine as Method 4.

15 ¹H nmr (DMSO) 8.25 (1H, s); 8.20 (1H,d); 7.95 (1H, d); 7.75 (3H,m); 7.35 (5H, m); 6.90 (2H,d); 5.35 (1H, m); 3.80 (3H,s); 3.70 (2H,m). MS TOF 417 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt
20 10.31 min.

Example 179.

R-2-{[N-(3-Amidino)benzoyl]amino}-2-phenylethyl-4-methylbenzamide

25 ¹H nmr (CD₃CN) 8.05 (1H, s); 8.00 (1H,d); 7.85 (1H, d); 7.65 (1H, t); 7.55 (2H,d); 7.35 (5H, m); 7.20 (2H,d); 5.30 (1H, m); 3.75 (2H,m); 2.30 (3H,s); MS TOF 401 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
30 water/acetonitrile/TFA) rt 10.58 min.

- 83 -

Compounds synthesised by Miscellaneous Methods

Example 180.

3-Amidino-benzoyl-DL-(4-methoxycarbonyl)phenylglycine-4-methylbenzylamine

D,L-4-Carboxyphenylglycine was prepared from 4-cyanobenzaldehyde by a method described by Clark, B. P. et al *Biorganic & Medicinal Chemistry Letters*, 1997, 7, 2777-2780.

D,L-4-Carboxyphenylglycine (600mg.) was suspended in methanol (30ml) at 0°C, saturated with hydrogen chloride gas and left at room temperature overnight. The reaction mixture was then evaporated to dryness under reduced pressure, resuspended in methanol and evaporated twice more and the solid triturated with ether, filtered and dried to give D,L-4-methoxycarbonylphenylglycine methyl ester HCl salt 620mg.

D,L-4-methoxycarbonylphenylglycine methyl ester HCl salt (600mg) was reacted with di-t-butyl dicarbonate in dioxane/ aqueous sodium bicarbonate to give after extraction with ethyl acetate and flash chromatography (SiO₂, hexane/ethyl acetate) - D,L-Boc-4-methoxycarbonylphenylglycine methyl ester 590mg. - ¹H nmr (CDCl₃) 7.78 (2H, d); 7.20 (2H, d); 5.61 (1H, broad d); 5.13 (1H, broad d); 3.63 (3H, s); 3.44 (3H, s); 1.16 (9H, s).

D,L-Boc-4-methoxycarbonylphenylglycine methyl ester (400mg) was dissolved in THF 10ml and treated with a solution of lithium hydroxide monohydrate 64mg. in water 1ml and the mixture stirred at room temperature for 30min. The reaction mixture was acidified with 2M hydrochloric acid, extracted with ethyl acetate and the organic solution was washed twice with water and dried

- 84 -

(MgSO₄). Evaporation of the solvent and purification by flash chromatography gave D,L-Boc-4-methoxycarbonyl-phenylglycine 282mg. ¹H nmr (CDCl₃) 8.0 (1H, broad d); 7.78 (2H, d); 7.20 (2H, d); 5.13 (1H, broad d); 3.80 (3H, s); 1.13 (9H, s).

D,L-Boc-4-methoxycarbonylphenylglycine was coupled to 4-methylphenyl-benzylamine using TBTU/DIPEA/DMF in the usual manner to give D,L-(4-methoxycarbonyl)phenylglycine-4-methylphenylbenzylamide. ¹H nmr (CDCl₃) 7.93 (2H, d); 7.35 (2H, d); 7.02 (2H, d); 6.93 (2H, d); 5.78 (2H, m); 5.08 (1H, broad s); 4.30 (2H, m); 3.86 (3H, s); 2.22 (3H, s); 1.30 (9H, s).

D,L-(4-methoxycarbonyl)phenylglycine-4-methylphenylbenzyl amide was deprotected using TFA/DCM and coupled to m-benzamidino carboxylic acid TFA salt as Method 4

¹H nmr (DMSO) 8.40 (1H, s); 8.30 (1H, d); 8.10 (3H, m); 7.80 (3H, m); 7.20 (4H, s); 6.00 (1H, s); 4.35 (2H, s); 4.00 (3H, s); 2.40 (3H, s); MS TOF 459 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA) rt 10.88 min.

25

Example 181.**3-(Trimethylacetyloxomethylcarbonylamidino)-benzoyl-D-phenylglycine-4-(4-chlorophenyl)piperazinamide**

Trimethylacetyloxymethyl carbonochloridate was prepared by the Method described by Folkmann and Lund, *Synthesis*, 1990, 1159., and reacted with 3-amidino-benzoyl-D-phenylglycine-4-(4'-chlorophenyl)piperazinamide (Example 116) in acetonitrile/aqueous sodium bicarbonate to give, after flash chromatography (SiO₂ - DCM/ethylacetate) trimethylacetyloxymethylcarbonyl-3-amidino-benzoyl-D-phenylglycine-4-(4'-chlorophenyl)piperazinamide.

- 85 -

¹H nmr (DMSO) 8.35 (1H, s); 8.10 (1H, d); 8.00 (1H, d); 7.60 (1H, t); 7.55 (2H, m); 7.40 (3H, m); 7.20 (2H, d); 6.80 (2H, d); 6.05 (1H, s); 5.75 (2H, s); 3.75 (1H, m); 3.65 (2H, m); 3.45 (3H, m); 3.30 (1H, m); 2.85 (1H, m) 1.20 (9H, s). MS TOF no (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 13.30 min.

Example 182.

R-3-Amidino-{2-(4-benzylpiperidin-1-yl)}-1-phenylethyl benzamide

Boc-D-Phenylglycine-OH was coupled to 4-benzylpiperidine as Method 4 and the Boc group removed with TFA. The product was dissolved in dry THF and added dropwise to LiAlH₄ (1.2eq) suspended in dry THF (15ml) under nitrogen and refluxed under nitrogen for 20 hours. On cooling the reaction was carefully diluted with water, acidified to pH 1 (HCl), and washed with EtOAc. The aqueous solution was basified to pH 10 (solid Na₂CO₃), then extracted with EtOAc to obtain product which was further elaborated as Method 4.

¹H nmr (CD₃CN) 8.25 (1H, s); 8.15 (1H, d); 7.95 (1H, d); 7.70 (1H, t); 7.55 - 7.10 (10H, m); 5.60 (1H, m); 3.70 (2H, m); 3.30 (2H, m); 2.95 (2H, m); 2.55 (2H, m); 1.80 (3H, m); 1.50 (2H, m). MS TOF 441 (M+1⁺). Hplc (Jupiter5 C18, Gradient 1, water/acetonitrile/TFA) rt 13.34 min.

Example 183.

3-(Benzoyloxycarbonylamidino)-benzoyl-D-phenylglycine-4-(4-chlorophenyl)piperazinamide

Example 116 was dissolved in DCM and DIPEA (2.2eq), then benzyl chloroformate (1.1eq) was added and stirred at room temperature for 45 mins. The reaction mixture was washed with water then chromatographed on silica gel 60 eluting with 65% EtOAc/Hexane to give the pure product.

- 86 -

¹H nmr (CD₃OD) 8.25 (1H, s); 8.05 (1H, d); 7.95 (1H, d);
7.60 (1H, t); 7.45 (5H, m); 7.20 (2H, d); 6.80 (2H, d);
6.10 (1H, s); 5.20 (2H, s); 3.80 (1H, m); 3.60 (3H, m);
3.20 (2H, m); 2.95 (1H, m); 2.50 (1H, m). MS TOF 611
5 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3,
water/acetonitrile/TFA) rt 13.15 min.

Example 184.

3-(tert-Butyloxycarbonylamidino)-benzoyl-D-
10 phenylglycine-4-(4-chlorophenyl)piperazinamide

Example 116 was dissolved in dioxane and NaHCO₃
(2.2eq.), di tertbutyl dicarbonate (1.1eq) was added and
stirred at room temperature for 1 week. The reaction
15 mixture was extracted with ethyl acetate and evaporated
to dryness. The product was chromatographed on silica
gel 60 eluting with 75% EtOAc/Hexane to give the pure
product.

20 ¹H nmr (CD₃OD) 8.25 (1H, s); 7.95 (2H, m); 7.55 (1H, t);
7.45 (5H, m); 7.15 (2H, d); 6.80 (2H, d); 6.10 (1H, s);
3.85 (3H, m); 3.60 (1H, m); 3.10 (2H, m); 2.90 (1H, m); 2.50
(1H, m); 1.45 (9H, s). MS TOF 577 (M+1⁺). Hplc (Jupiter5
C18, Gradient 1, water/acetonitrile/TFA) rt 12.72 min.

25

Example 185.

3-(Acetylamidino)-benzoyl-D-phenylglycine-4-(4-
chlorophenyl)piperazinamide

30 Example 116 was dissolved in dichloromethane and DIPEA
(2.2eq) added. Acetyl chloride (1.1eq) was added and
stirred at room temp for 45 mins. The reaction mixture
was then washed with water and chromatographed on
silica gel 60 eluting with EtOAc. The product was
35 further purified by preparative hplc.

¹H nmr (CD₃OD) 8.25 (1H, s); 8.05 (2H, m); 7.65 (1H, t);

- 87 -

7.45 (5H, m); 7.25 (2H, d); 6.85 (2H, d); ; 6.15 (1H, s);
3.65 (3H, m); 3.60 (1H, m); 3.20 (2H, m); 3.00 (1H, m); 2.55
(1H, m); 2.40 (3H, s). MS TOF 519 (M+1⁺). Hplc (Jupiter5
C18, Gradient 3, water/acetonitrile/TFA) rt 14.53 min.

5.

Example 186.**(R)-O-(3-Amidinobenzyl)-N-(4-methylbenzyl)mandelamide****(R)-O-(3-Cyanobenzyl) mandelic acid**

10 (R)-Mandelic acid 91g, 6.57mmol.) was dissolved in THF
(15ml.) and treated portionwise with sodium hydride
(60%w/w, 0.798g, 19.7mmol.). The white suspension was
heated at reflux and treated dropwise with 3-
15 cyanobenzylbromide (1.93g, 9.86mmol.) in THF (20ml.)
over 5mins. The mixture was refluxed for 2 days. The
reaction mixture was cooled and treated with water
(100ml.). The aqueous phase was washed with diethyl
ether then acidified to pH 1 with 10% HCl solution. The
20 product was extracted with ethyl acetate. The extracts
were washed with brine and dried over magnesium
sulphate, filtered and evaporated to give the product
(1.5g) as a gum.

¹H nmr (CD₃OD) 7.70 (3H, m); 7.40 (6H, m); 5.00 (1H, s);
25 4.65 (2H, m).

(R)-O-(3-Cyanobenzyl)-N-(4-methylbenzyl)mandelamide

HOAt (0.27g, 1.96mmol.) and DBU (0.3g, 1.96mmol.) was
30 dissolved in DMF (8ml.) and cooled in ice. EDC (0.38g,
1.96mmol.) was added in 3 portions and stirred for
10mins.. To this mixture was added (R)-O-(3-Cyanobenzyl)
mandelic acid (0.5g, 1.87mmol.) in DMF (2ml.) and the
mixture allowed to warm to room temperature for 5 mins.
35 4-Methylbenzylamine (0.23g, 1.96mmol.) was then added in
one portion. The mixture was stirred for 1 hr. The
reaction was quenched with water (100ml.) and basified

- 88 -

with 2M sodium hydroxide solution. The product was extracted with ethylacetate. The combined extracts were washed with dilute (5%) lithium bromide solution, water and brine, dried over magnesium sulphate, filtered and evaporated to give a gum (0.58g). The crude product was chromatographed on silica gel eluting with 35% ethylacetate in hexane.

¹H nmr (CDCl₃) 7.50 (2H,m); 7.30 (6H,m); 7.05 (4H,s); 6.90 (1H,m); 4.80 (1H,s); 4.55 (2H,m) 4.35 (2H,s); 2.25 (3H,s).

(R)-O-(3-Amidinobenzyl)-N-(4-methylbenzyl)mandelamide

(R)-O-(3-Cyanobenzyl)-N-(4-methylbenzyl)mandelamide (0.16g) was dissolved in ethanol (10ml) and treated with hydrogen chloride gas for 30 mins. The mixture was then stirred overnight at room temperature, evaporated to dryness and dissolved again in ethanol (10ml.). The mixture was treated with ammonia gas for 5 mins. and allowed to stand overnight. The mixture was evaporated and the product purified by preparative hplc.

¹H nmr (CDCl₃) 7.65 (1H,s); 7.50 (1H,d); 7.30 (7H,m); 7.00 (4H,s); 4.80 (1H,s); 4.55 (2H,m) 4.35 (2H,s); 2.25 (3H,s). MS TOF 387 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 11.24 min.

Example 187.

2-(3-Amidinobenzoyl)-1-(3-phenylpropanoyl)-1-phenylhydrazine.

1-(3-phenylpropanoyl)-1-phenylhydrazine

2-tert Butoxycarbonyl-1-phenylhydrazine (1.3g) was dissolved in pyridine (6ml.) and treated with 3-phenylpropanoyl chloride (0.75ml.). The mixture was stirred overnight at room temperature under argon. The pyridine was then removed by evaporation and the residue

- 89 -

taken up in ethyl acetate and washed with 10% hydrochloric acid, saturated sodium carbonate solution and brine. The ethyl acetate fraction was dried over magnesium sulphate, filtered and evaporated. The crude product was chromatographed on silica eluting 5-25% ethyl acetate in hexane. The purified product was dissolved in DCM and treated with TFA before coupling to m-benzamidine by Method 4.

¹H nmr (CD₃OD) 8.20 (1H, s); 8.05 (1H,d); 7.90 (1H,d); 7.65 (1H, t); 7.35 (5H, m); 7.10 (5H, m); 6.15 (1H, s); 2.60 (4H,m). MS TOF 388 (M+1⁺). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 10.75 min.

Example 188.

3-{[R]-N-(3-Amidinobenzamido)-1-phenyl}methyl-5-benzyl-1,2,4-triazole.

EDC (1.6g) was dissolved in DMF (20ml.) and treated with HOAt (1.16g) in DMF (10ml.) and stirred for 10 mins. DIPEA (1.46ml.) was then added and the stirring continued for a further 15 mins. Boc-D-phenylglycine (2g) in DMF (10ml.) was then added dropwise and stirring continued for a further 25mins. 1M Hydrazine in THF (84ml.) was then added and the mixture stirred over night. The solvents were then evaporated and the residue treated with water (100ml.), basified with 2M sodium hydroxide solution and extracted with ethyl acetate. The combined extracts were washed with water and brine, dried with magnesium sulphate, filtered and evaporated to give a yellow solid (1.38g).

¹H nmr (CDCl₃) 7.20 (5H,s); 5.55 (1H,s); 1.25 (9H,s).

The above product (0.3g, 1.13mmol.) was dissolved in ethanol (6ml.) and added dropwise to a solution of benzyl ethyl imidate hydrochloride in ethanol (3ml.) at

- 90 -

room temperature. The mixture was stirred for 30 mins. The solvent was then evaporated and the residue treated with dilute sodium bicarbonate solution and extracted with ethylacetate. The combined extracts were washed
5 with brine, dried over magnesium sulphate, filtered and evaporated to give a yellow gum. (0.43g).

^1H nmr (CDCl_3) 7.30 (10H,m); 5.80 (1H,s); 3.50 (2H,m);
1.45 (9H,s).

10

The above crude product (0.43g, 1.12 mmol.) was dissolved in xylene (60ml.) with para toluenesulphonic acid (cat.) and heated to 160°C for 1 day. The xylene was removed under vacuo and the triazole (0.35g) used
15 without further purification.

^1H nmr (CDCl_3) 7.20 (10H,m); 5.70 (1H,s); 4.00 (2H,m);
1.30 (9H,s).

20

The above product was treated with TFA and coupled to m-benzamidine by Method 4

^1H nmr (CD_3OD) 8.20 (1H,s); 8.05 (1H,d); 7.80 (1H,d);
7.60 (1H,t); 7.20 (10H,m); 6.50 (1H,s); 4.10 (2H,s).

25

MS TOF 411 ($\text{M}+1^+$). Hplc (Jupiter5 C18, Gradient 3, water/acetonitrile/TFA) rt 14.38 min.

References:

30

(1)(a) Bucherer, H. T.; Steiner W. *J. Prakt. Chem* 1934, 140, 291-316.

(b) Greene T. W.; Wuts P. G. M. *Protective Groups in Organic Synthesis*, (John Wiley and Sons, Inc. 1991) 318-319 and references therein.

35

(c) Bolin D. R.; Sytwu I.; Humiec F.; Meienhofer J. *Int. J. Peptide Protein Res.* 1989, 33, 353-359.

(2) Shearer, B. G., Oplinger, J. A., Lee, S. *Tet. Letts.* 38(2), 179-182 (1997)

(3) Chandrakumar, N.S. *Synthetic Comms.*, 26(14), 2613-2616 (1996)

5 (4) Bundle et al *J. Chem, Soc. Perkin Trans I* 1985 2247)

Assay protocols

Enzyme Inhibition assays:

10

Enzyme assays were carried out at room temperature in 0.1M phosphate buffer, pH7.4 according to the method of Tapparelli et al (*J. Biol. Chem.* 1993, 268, 4734-4741).

15 Purified human factor Xa, trypsin, thrombin and plasmin were purchased from Alexis Corporation, Nottingham, UK. Urokinase was purchased from Calbiochem, Nottingham, UK. Chromogenic substrates for these enzymes;

20 pefachrome-FXA, pefachrome-TRY, pefachrome-TH, pefachrome-PL and pefachrome-UK were purchased from Pentapharm AG, Basel, Switzerland. Product (*p*-nitroaniline) was quantified by adsorption at 405nm in 96 well microplates using a Dynatech MR5000 reader (Dynex Ltd, Billingshurst, UK). K_m and K_i were calculated using SAS PROC NLIN (SAS Institute, Cary, NC, USA, Release 6.11) K_m values were determined as 25 100.9 μ M for factor Xa/pefachrome-FXA and 81.6 μ M for trypsin/pefachrome-TRY. Inhibitor stock solutions were prepared at 40mM in Me2SO and tested at 500 μ M, 50 μ M and 5 μ M. Accuracy of K_i measurements was confirmed by 30 comparison with K_i values of known inhibitors of factor Xa and trypsin.

In agreement with published data, benzamidine inhibited factor Xa, trypsin, thrombin, plasmin and urokinase 35 with K_i values of 155 μ M, 21 μ M, 330nM, 200nM and 100nM respectively. NAPAP inhibited thrombin with a K_i value of 3nM. Compounds of the invention were found to have activity in these assays.

Antithrombotic Activity

The test material (Factor Xa inhibitor) was administered either intravenously, intraperitoneally or orally to a group of rats for experiment. A second group received vehicle (saline) only as a control, and a third group of animals received a standard antithrombotic (subcutaneous lmw heparin) as a positive control.

10

To perform the experiment male Sprague-Dawley rats (250-400g in weight) were anaesthetised by the inhalation of isoflurane with the addition of oxygen and nitrous oxide. The left or right femoral vein was carefully exposed and isolated from the femoral artery and saphenous nerve. Following removal of connective tissue a cannula containing physiological saline was inserted into the femoral vein.

20

A segment of each of the left and right jugular vein was exposed and separated from the surrounding connective tissue. Each segment consisted of the section of vein between the exit point from the thorax to the vessel's first major junction.

25

At the desired interval following the administration of the test material or vehicle a bolus injection of 'deactivated' human serum (1.32 ml.kg^{-1}) was administered over less than 30 seconds, via the femoral vein cannula. Two minutes following the thrombus challenge both the jugular vein segments were ligatured at both ends and left *in situ* for thrombi to form.

After 10 min both jugular segments were carefully excised, and placed in a petri dish containing 0.9% saline. A blood sample (1.8 ml blood + 0.2 ml 3.8% sodium citrate) was obtained by cardiac puncture and the animal sacrificed by an overdose of Expiral (sodium

35

pentobarbitone) administered intravenously via the femoral vein cannula or by cardiac puncture. The 2 segments of jugular vein were carefully dissected longitudinally along one surface to expose the lumen and dispel the vessel contents into the saline. The tissues were examined for the presence of any developed thrombi and scored accordingly.

Thrombus score:

10

- 0 = No thrombus
- 1 = One or several small thrombi
- 2 = Several larger thrombi
- 3 = Large thrombus occluding the vessel

15

Compounds of the invention were found to have significant antithrombotic activity in these assays.

Partial Thromboplastin Time Test Protocol

20

Venous blood was collected into 3.2% (0.109M) trisodium citrate vacutainer tubes at 1 volume of anticoagulant to nine volumes of blood.

25

The blood cells were separated by centrifugation at 700g for ten minutes to yield plasma, which was frozen at 70°C until required.

30

To perform the test, 100µl of plasma was pipetted into a glass test tube, 1µl of test compound in DMSO was added, and allowed to warm to 37° over two minutes.

35

100µl of warm (37°) Manchester (tissue thromboplastin) reagent (Helena Biosciences, UK) was added, allowed to equilibrate for sixty seconds.

100µl of warm (37°) 25mM calcium chloride solution was added to initiate clotting.

The test tube was tilted three times through a 90° angle every five seconds to mix the reagents and the time to clot formation recorded.

- 5 Data from a series of observations and test compound concentrations are analysed by a SAS statistical analysis program and a CT2 (Concentration required to double clotting time) for each compound is generated.
- 10 Compounds of the invention were found to significantly elongate the partial thromboplastin time.

Kaolin Activated Partial Thromboplastin Time Test Protocol

- 15 Venous blood was collected into 3.2% (0.109m) trisodium citrate vacutainer tubes at 1 volume of anticoagulant to nine volumes of blood.
- 20 The blood cells were separated by centrifugation at 700g for ten minutes to yield plasma, which was frozen at 70°C until required.
- 25 To perform the test, 100µl of plasma was pipetted into in a glass test tube, 1µl of test compound in DMSO was added, and allowed to warm to 37° over two minutes.
- 30 100µl of APTT reagent resuspended in Owren's buffer (Thrombosis Reference Centre University of Manchester, Withington Hospital, Manchester) was added, mixed thoroughly, and 100µl of prewarmed kaolin suspension in Owren's buffer added.
- 35 This mixture was incubated for ten minutes at 37°, with regular gentle mixing by tilting through a 90° angle, three times sequentially at one minute intervals to maintain the kaolin in suspension.

After ten minutes 100 μ l of 25mM calcium chloride solution was added, the suspension redispersed by mixing as above and the time to clot formation recorded.

5

Data from a series of observations and test compound concentrations are analysed by a SAS statistical analysis program and a CT2 (Concentration required to double clotting time) for each compound is generated.

10

Table 1 Biological data for selected examples


15	EX No	FXa pKi	thrombin pKi	trypsin pKi	APTT CT2 uM	PT CT2 uM
	84	7.98	5.04	5.61	5.8	1.9
	89	7.85	5.99	6.61	NT	NT
	63	7.80	5.96	6.54	NT	3.7
	87	7.62	5.10	5.89	14	4.2
20	33	7.50	6.14	6.39	NT	7.1
	32	7.39	6.09	5.65	7.4	5.6
	65	7.19	5.93	5.79	9.7	13
	36	7.12	6.24	6.16	9.4	9
	64	7.06	5.72	5.91	7.6	8.3
25	68	7.01	5.32	5.59	NT	17
	66	6.94	6.66	6.55	4.5	4.6
	31	6.80	6.98	5.56	3.7	3.2
	37	6.79	4.77	5.84	36	15
	80	6.57	5.16	5.73	NT	13
30	12	6.39	5.27	6.10	6.5	3.4

Claims

1. A serine protease inhibitor compound of formula I

5

10



(where R_1 and R_2 each independently is hydrogen or hydroxyl, alkoxy, alkyl, aminoalkyl, hydroxyalkyl alkoxyalkyl, alkoxycarbonyl, acyloxymethoxycarbonyl or alkylamino optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl;

each R_3 independently is R_1 , R_2 , amino, halo, cyano, nitro, thiol, alkylthio, alkylsulphonyl, alkylsulphenyl, alkylsulphonamido, alkylaminosulphonyl, haloalkoxy and haloalkyl;

each X independently is a C, N, O or S atom or a CO, CR_1 , $C(R_1)_2$ or NR_1 group, at least one X being C, CO, CR_1 or $C(R_1)_2$, with the proviso that if the benzamidine group is unsubstituted and the X-X group is $-CH_2C(R_1)_2-$ then R_1 is hydrogen or is attached to the alkylene carbon atom by a heteroatom;

L is an organic linker group containing 1 to 5 backbone atoms selected from C, N, O and S, or a branched alkyl or cyclic group;

Y is a nitrogen atom or a CR_1 group or Y and L taken together form a cyclic group;

Cy is a saturated or unsaturated, mono or poly cyclic, homo or heterocyclic group optionally substituted by groups R_3 or phenyl optionally substituted by R_3 ;

Lp is a lipophilic organic group, selected from an

- 97 -

alkyl, heterocyclic, alkenyl, alkaryl, cycloalkyl, polycycloalkyl, cycloalkenyl, aryl, aralkyl or haloalkyl group or a combination of two or more such groups optionally substituted by one or more of oxa, oxo, aza, thio, halo, amino, hydroxy or R_3 groups;

D is a hydrogen bond donor group; and n is 0, 1 or 2)

or a physiologically tolerable salt thereof.

2. A compound as claimed in claim 1 wherein R_1 represents hydrogen.

3. A compound as claimed in either one of claims 1 or 2 wherein R_2 is hydrogen, alkoxycarbonyl or hydroxy.

4. A compound as claimed in any one of claims 1 to 3 wherein R_3 is hydrogen, amino, aminoalkyl, alkylamino, hydroxy, hydroxyalkyl, thiol, alkylthio or alkoxy.

5. A compound as claimed in claim 4 wherein R_3 is para to the amidino group.

6. A compound as claimed in claim 5 wherein R_3 is amino.

7. A compounds as claimed in any one of claims 1 to 6 wherein the linker X-X is -CH=CH-, -CONH-, -CONR₁-, -NH-CO-, -NH-CH₂-, -CH₂-NH-, -CH₂O-, -OCH₂-, -COO-, -OC=O- or -CH₂CH₂-.

8. A compound as claimed in claim 7 wherein the X moiety nearest Y is NH.

9. A compound as claimed in claim 7 wherein the X moiety nearest the phenyl ring is CH₂ or CO.

- 98 -

10. A compound as claimed in claim 7 wherein X-X is
-CONH-.
11. A compound as claimed in any one of claims 1 to 10
5 wherein Y is CH.
12. A compound as claimed in any one of claims 1 to 11
wherein Cy represents aryl or heteroaryl optionally
substituted by R_3 .
- 10 13. A compound as claimed in claim 12 wherein R_3
represents phenyl optionally substituted by R_3 or naphthyl
optionally substituted by R_3 .
- 15 14. A compound as claimed in claim 13 wherein Cy
represents phenyl.
15. A compounds as claimed in any one of claims 1 to 14
wherein the linker L represents CO, CH_2NH , $\text{CONR}_1(\text{CH}_2)_m$,
20 $(\text{CH}_2)_m\text{N}(\text{R}_1)\text{CO}(\text{CH}_2)_m$, $(\text{CH}_2)_{m+2}$, $(\text{CH}_2)_m\text{CO}(\text{CH}_2)_m$, $(\text{CH}_2)_m\text{OC}=\text{O}$,
 $(\text{CH}_2)_m\text{O}$ or $\text{CH}=\text{CH}(\text{CH}_2)_m$ (where each m is independently 0 or
1).
16. A compound as claimed in claim 15 wherein the
25 linker L represents $\text{CONR}_1(\text{CH}_2)_m$, CO or $\text{CH}_2\text{NR}_1\text{CO}$ where m is
0 or 1.
17. A compound as claimed in any one of claims 1 to 16
wherein said lipophilic group comprises cycloalkyl,
30 azacycloalkyl, diazacycloalkyl, phenyl, naphthyl,
adamantyl, decalinyl, tetrahydrodecalinyl, bicycloalkyl,
mono- or diazabicycloalkyl, mono- or bicyclo
heteroaromatic or a linear or branched alkyl, alkylene,
alkenyl or alkenylene group all optionally substituted
35 by one or more groups R_3 , or a combination of at least
two such groups linked by a spiro linkage or a single or
double bond or by C=O, O, S, SO, SO₂, CONR_1 , $\text{NR}_1\text{-CO-}$, NR_1

- 99 -

linkage.

18. A compound as claimed in claim 17 wherein the lipophilic groups is a methyl-cyclohexyl, methylcyclohexylmethyl, methylphenylmethyl, phenylethyl, benzylpiperidinyl, benzoylpiperidinyl, bispiperidinyl or phenylpiperazinyl.

19. A compound as claimed in any one of claims 1 to 18 wherein the hydrogen bond donor group is attached to the lipophilic group.

20. A compound as claimed in claim 19 wherein nitrogen or oxygen is the hydrogen bond donor atom.

21. A compound as claimed in claim 20 wherein the hydrogen bond donor is formed from a hydroxyl group, a primary, secondary or tertiary amine, or a primary or secondary imine group as part of an amidine or guanidine or a saturated or unsaturated heterocyclic group containing 5 to 7 ring atoms and having a ring nitrogen.

22. A compound as claimed in any one of claims 1 to 21 wherein Y is carbon and preferably has the conformation that would result from construction from a D- α -aminoacid $\text{NH}_2\text{-CR}_1(\text{Cy})\text{-COOH}$.

23. A compound as claimed in any one of claims 1 to 22 wherein the serine protease to be inhibited is trypsin, thrombin, urokinase or Factor Xa.

24. A compound as claimed in claim 1 being 3-Amidino-benzoyl-D-phenylglycine-4-benzoyl piperidinamide.

25. A pharmaceutical composition comprising a serine protease inhibitor as claimed in any one of claims 1 to 24 together with at least one pharmaceutically

- 100 -

acceptable carrier or excipient.

26. A serine protease inhibitor as claimed in any one
of claims 1 to 24 for use in combatting a condition
5 responsive to said inhibitor.

27. Use of a serine protease inhibitor as claimed in
any one of claims 1 to 24 for the manufacture of a
medicament for use in a method of treatment of the human
10 or non-human animal body to combat a condition
responsive to said inhibitor.

28. A compound as claimed in claim 27 wherein said
condition is a thrombotic disease, asthma, emphysema,
15 cirrhosis, arthritis, carcinoma, melanoma, restenosis,
atheroma, trauma, shock and reperfusion injury.

29. A method of treatment of the human or non-human
animal body to combat a condition responsive to a serine
20 protease inhibitor, said method comprising administering
to said body an effective amount of a serine protease
inhibitor as claimed in any one of claims 1 to 24.

INTERNATIONAL SEARCH REPORT

In. Application No
PCT/GB 98/02605

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C07K5/065 C07K5/087 C07K5/107 A61K38/05 A61K38/06 A61K38/07 C07C257/18 C07D209/20 C07D317/60 C07D333/24 C07D207/14 C07D211/26 C07D207/16 C07D233/54 C07D211/30		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C07K A61K C07C C07D		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CHEMICAL ABSTRACTS, vol. 103, no. 25, 23 December 1985 Columbus, Ohio, US; abstract no. 209699z, XP002083330 see abstract & R. FERRONI ET AL.: FARMACO, ED. SCI., vol. 40, no. 10, 1985, pages 717-729, ---	1-28
Y	CHEMICAL ABSTRACTS, vol. 108, no. 7, 15 February 1988 Columbus, Ohio, US; abstract no. 48724q, XP002083331 see abstract & R. FERRONI ET AL.: FARMACO, ED. SCI., vol. 42, no. 10, 1987, pages 709-715, --- -/--	1-28
<div style="display: flex; justify-content: space-between;"> <input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex. </div>		
* Special categories of cited documents : <div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-weight: bold;">6 November 1998</div>		Date of mailing of the international search report <div style="text-align: center; font-weight: bold;">24/11/1998</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer <div style="text-align: center; font-weight: bold;">Frelon, D</div>

INTERNATIONAL SEARCH REPORT

onal Application No

PCT/GB 98/02605

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D211/22 C07D211/34 C07D217/26 C07D295/18 C07D295/22
C07D277/28 C07D249/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CHEMICAL ABSTRACTS, vol. 71, no. 7, 18 August 1969 Columbus, Ohio, US; abstract no. 28982m, XP002083332 see abstract & B.R. BAKER ET AL.: JOURNAL OF MEDICINAL CHEMISTRY, vol. 12, no. 3, 1969, pages 408-414, WASHINGTON US	1-28
Y	EP 0 672 658 A (LILLY CO ELI) 20 September 1995 see examples 16,24,25,27,31,33,36,37,40,41 --- -/--	1-28

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier document but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
"&" document member of the same patent family

Date of the actual completion of the international search

6 November 1998

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Frelon, D

INTERNATIONAL SEARCH REPORT

Int. Application No

PCT/GB 98/02605

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 97 24135 A (LU TIANBAO ;TOMCZUK BRUCE E (US); ILLIG CARL (US); SOLL RICHARD M) 10 July 1997 see examples 6,7,15,17 ---	1-28
Y	WO 95 13274 A (PFIZER LTD ;PFIZER RES & DEV (IE); PFIZER (US); DANILEWICZ JOHN CH) 18 May 1995 see claims ---	1-28
Y	WO 92 08709 A (PENTAPHARM AG) 29 May 1992 see claims ---	1-28
A	EP 0 483 667 A (THOMAE GMBH DR K) 6 May 1992 see ex.47, page 82; ex.140, page 88; ex. 136, page 103; page 23 ---	1-28
A	WO 94 29336 A (ASTRA AKTIEBOLAG) 22 December 1994 see abstract ---	1-28
A,P	WO 97 36859 A (RICO JOSEPH G ;SEARLE & CO (US); YU STELLA S (US); CHEN BARBARA B) 9 October 1997 see examples 13-20 ---	1-28
A	EP 0 588 655 A (ONO PHARMACEUTICAL CO) 23 March 1994 see abstract ---	1-28
A	CHEMICAL ABSTRACTS, vol. 101, no. 1, 2 July 1984 Columbus, Ohio, US; abstract no. 6764c, XP002083333 see abstract & H. VIEWEG ET AL.: PHARMAZIE, vol. 38, no. 12, 1983, pages 818-820, BERLIN DE -----	1-28

INTERNATIONAL SEARCH REPORT

.national application No.

PCT/GB 98/ 02605

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 29
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim 29
is directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims: it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

In view of the extremely large scope claimed through unspecific and inconcise expressions/notations like "organic linker", "cyclic group", "lipophilic group", "alkyl", "heterocyclic group" and the like, it is not possible to fully and clearly specify the matter for which protection is sought. This contravenes the PCT requirements (e.g., clarity and conciseness, Art. 6 PCT).

In other words, the scope is so large and the portion which is always present is so small that it is not possible to carry out a complete and meaningful search into the state of the art on the basis of the claims. The number of theoretically conceivable compounds resulting from the combination of all claimed substituents precludes a comprehensive search. "No special search effort need to be made for searching unduly wide or speculative claims" (Guidelines B-III, 3.7 and 3.12) and the costs of the search are "to be kept within reasonable bounds" (Guidelines B-III, 2.1).

Furthermore, many dissimilar structures are encompassed for which no evidence is in the description that they all are solution of the problem underlying the application, namely the provision of compounds inhibiting serine protease. As a matter of principle, a formula consisting virtually completely of variables is hardly a permissible generalisation which is fairly based on experimental evidence. All potentially novel and inventive subject-matter is hidden in this large amount of possible compounds (Art. 33 PCT).

The search has been limited to the subject-matter as illustrated by the prepared compounds for which the nomenclature is clear (Art. 17 PCT) and to the general idea (serine protease inhibitors) underlying the application.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 98/02605

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0672658	A	20-09-1995	AU 684918 B	08-01-1998
			AU 1975295 A	18-09-1995
			BR 9506979 A	18-11-1997
			CA 2183464 A	09-08-1995
			CN 1147205 A	09-04-1997
			CZ 9602584 A	11-06-1997
			FI 963451 A	03-09-1996
			HU 76330 A	28-08-1997
			JP 9509937 T	07-10-1997
			NO 963684 A	28-10-1996
			NZ 282588 A	19-12-1997
			PL 320637 A	13-10-1997
			WO 9523609 A	08-09-1995
			US 5705487 A	06-01-1998
			US 5726159 A	10-03-1998
			US 5707966 A	13-01-1998
WO 9724135	A	10-07-1997	AU 1432397 A	28-07-1997
			NO 982987 A	28-08-1998
WO 9513274	A	18-05-1995	AT 155464 T	15-08-1997
			AU 680565 B	31-07-1997
			AU 8104994 A	29-05-1995
			CA 2176037 A	18-05-1995
			CN 1134148 A	23-10-1996
			CZ 9601322 A	14-08-1996
			DE 69404325 D	21-08-1997
			DE 69404325 T	30-10-1997
			DK 728132 T	01-09-1997
			EP 0728132 A	28-08-1996
			ES 2105868 T	16-10-1997
			FI 961939 A	07-05-1996
			GR 3024228 T	31-10-1997
			HU 76268 A	28-07-1997
			JP 2739776 B	15-04-1998
			JP 9500391 T	14-01-1997
			NO 961841 A	07-05-1996
			NZ 275798 A	22-09-1997
			PL 314237 A	02-09-1996
			US 5750520 A	12-05-1998

INTERNATIONAL SEARCH REPORT

Information on patent family members

In International Application No

PCT/GB 98/02605

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9513274 A		ZA 9408772 A	07-05-1996
WO 9208709 A	29-05-1992	AU 8868991 A	11-06-1992
		CA 2073776 A	16-05-1992
		EP 0511347 A	04-11-1992
		JP 5503300 T	03-06-1993
		US 5518735 A	21-05-1996
EP 0483667 A	06-05-1992	DE 4035961 A	07-05-1992
		AT 163008 T	15-02-1998
		AU 650488 B	23-06-1994
		AU 8692691 A	07-05-1992
		CA 2054850 A	03-05-1992
		DE 59108930 D	12-03-1998
		ES 2113867 T	16-05-1998
		FI 915136 A	03-05-1992
		HU 67288 A	28-03-1995
		HU 9500156 A	28-12-1995
		IL 99926 A	18-06-1996
		JP 4264068 A	18-09-1992
		MX 9203677 A	01-07-1992
		NO 174806 B	05-04-1994
		NZ 240463 A	24-04-1997
		PT 99402 A	30-09-1992
		RU 2040519 C	25-07-1995
		US 5541343 A	30-07-1996
		US 5591769 A	07-01-1997
		ZA 9108734 A	04-05-1993
WO 9429336 A	22-12-1994	AU 684086 B	04-12-1997
		AU 6986994 A	03-01-1995
		BR 9406746 A	19-03-1996
		CA 2162900 A	22-12-1994
		CN 1127509 A	24-07-1996
		CZ 9503020 A	17-04-1996
		EP 0701568 A	20-03-1996
		FI 955828 A	04-12-1995
		HR 940311 A	31-10-1996
		HU 74739 A	28-02-1997
		JP 8511018 T	19-11-1996

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 98/02605

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9429336 A		LT 1947 A, B	27-12-1994
		MX 9404114 A	31-01-1995
		NO 954873 A	01-02-1996
		NZ 267534 A	22-08-1997
		PL 311819 A	18-03-1996
		SG 48013 A	17-04-1998
		SK 145495 A	01-10-1996
		US 5780631 A	14-07-1998
		US 5602253 A	11-02-1997
		US 5783563 A	21-07-1998
		US 5723444 A	03-03-1998
WO 9736859 A	09-10-1997	AU 2536097 A	22-10-1997
EP 0588655 A	23-03-1994	AT 145894 T	15-12-1996
		CA 2106452 A	19-03-1994
		DE 69306345 D	16-01-1997
		DE 69306345 T	23-10-1997
		DK 588655 T	23-12-1996
		ES 2097457 T	01-04-1997
		GR 3022642 T	31-05-1997
		JP 2736967 B	08-04-1998
		JP 8259512 A	08-10-1996
		JP 2736952 B	08-04-1998
		JP 8109164 A	30-04-1996
		US 5432178 A	11-07-1995
		US 5622984 A	22-04-1997
		US 5614555 A	25-03-1997